


## Synthetic compounds useful as nodulation agents of leguminous plants and preparation processes thereof

### I. DESCRIPTION

5 The present invention relates to synthetic compounds that are active on plants, especially as legume nodulation factors, and also as plant growth stimulators, and to methods for preparing such compounds.

10 It is known that the process of nitrogen fixing by legumes is based on symbiosis between these plants and soil bacteria, the Rhizobia. The Rhizobium-legume symbiosis produces each year, by means of the root nodules, more ammonium than all of the nitrogen fertilizer industry. This symbiosis thus plays a considerable agronomic role. Legumes are very rich in proteins and produce about one third of the plant proteins consumed worldwide, by virtue of pulses such as soybean, pea, horse bean, groundnut, bean and lupin, and forage plants such as alfalfa and clover.

15 The formation of nitrogen-fixing nodules starts with an exchange of molecular signals, flavonoids secreted by the plant and nodulation factors (Nod factors) synthesized by the bacterium. These factors consist of an oligosaccharide fragment and a lipid chain attached to this skeleton on the non-reducing end. They have structural attributes (substitutions on the sugars at the two ends and variability of the chain) that make them specific to the legume-bacterium couple.



20

These lipochito-oligosaccharides (LCO) may be either isolated directly from a particular culture of rhizobia, synthesized chemically, or obtained chemo-enzymatically. Via the latter method, the oligosaccharide skeleton may be formed by culturing of recombinant *Escherichia coli* bacterial strains in a fermenter, and the lipid chain may then be attached chemically.

25


Treatment of the seeds of legumes, for instance soybean, with Nod factors at very low concentrations, may result in a large increase in the number of nitrogen-fixing root nodules and a significant increase in the yields, under agronomic conditions. It is thus clear that, in the future, compounds of Nod factor type will be produced industrially for large-scale agronomic use. However, the industrial preparation and conditioning of natural Nod factors presents two types of drawback: (1) the natural Nod factors are difficult

30

to assay via simple methods such as spectrometric methods; (2) they are unstable in the presence of plants or in soils, in particular because they have a -CO-NH- bond that may be broken by plant or microbial enzymes present in the rhizosphere.

One of the aspects of the present invention is to propose a process for preparing compounds of Nod factor type, some of these compounds constituting another aspect of the invention. Specifically, certain biologically active compounds show strong absorption in the ultraviolet range, which makes them easy to assay during their industrial preparation and allows them to be detected and assayed easily in the product intended for marketing, and allows their stability and storage in such products to be tested. In addition, some of these synthesized compounds show higher stability than the natural Nod factors.

The target Nod factor is especially the factor associated with alfalfa (**1**) as described above. Among all the existing legumes, alfalfa and vetch have been the subject of numerous studies. By virtue of several activity tests performed with various Nod factor analogs, it has been possible to establish a relationship between the structure of the LCO and the plant response. The isolated nodulation factor that is most active on alfalfa is a tetramer sulfated in position 6 of the reducing sugar, acetylated in position 6 of the sugar located at the nonreducing end and acylated with a chain of 16 carbons containing two unsaturations (C16:2 $\Delta$ 2E,9Z) (**1**)



It has been shown that the absence of acetate, the addition of a glucosamine unit, or the loss of one of the two unsaturations produce only a moderate reduction in activity. A test of variation of the chain length revealed maximum activity for a chain of 16 carbons. Finally, the sulfate is of prime importance for recognition of the Nod factor by alfalfa, but, on the other hand, its absence is necessary for action in vetch.


Studies have shown that the conjugated unsaturation present at 2 of the lipid chain was important for nodulation, since an analog acylated with a C16:1 $\Delta$ 9Z chain is less active (by just under a factor of 10).

From these results, a series of analogs for which the conjugated amide bond is mimicked by a benzamide bond have been prepared. These compounds constitute one of the aspects of the present invention.

Another series of analogs containing a function of benzylamine type constitute another aspect of the invention.

Such compounds are capable of causing or promoting the phenomenon of nodulation in legumes and of stimulating plant growth and development.

- 5 The compounds that are plant nodulation factors according to the invention are preferably compounds of formula (I)



10 in which

- ▶ n represents 1, 2 or 3;
- ▶ A represents a substituent chosen from -C(O)-, -C(S)-, -CH<sub>2</sub>-, -CHR<sup>10</sup>-, -CR<sup>10</sup>R<sup>11</sup>-, -C(O)O-, -C(O)S-, -C(S)O-, -C(S)S-, -C(O)NH-, -C(NH)NH- and -C(S)NH-;

- ▶ B represents

- an arylene;
- a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a naphthylene;
- a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a divalent radical derived from 2 fused aromatic rings of 5 or 6 atoms each;
- a divalent radical derived from 2 fused aromatic or heteroaromatic rings of 5 or 6 atoms each, comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a biphenylene;
- or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents R<sup>12</sup> and R<sup>13</sup> chosen, independently of each other, from halogen, CN, C(O)OR<sup>14</sup>, C(O)NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, OCF<sub>3</sub>, -NO<sub>2</sub>, N<sub>3</sub>, OR<sup>14</sup>, SR<sup>14</sup>, NR<sup>15</sup>R<sup>16</sup> and C<sub>1-6</sub>-alkyl;

- ▶ C represents a substituent chosen from -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>17</sup>-, -CR<sup>17</sup>R<sup>18</sup>- and -NR<sup>19</sup>;

- ▶ D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;

► E and G represent, independently of each other, a substituent chosen from H, OH, OR<sup>20</sup>, NH<sub>2</sub> and NHR<sup>20</sup>;

► R<sup>1</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, C(O)H and C(O)CH<sub>3</sub>;

► R<sup>2</sup>, R<sup>3</sup>, R<sup>6</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup> and R<sup>19</sup> represent, independently of each other, a substituent chosen from H, C<sub>1-6</sub>-alkyl, C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;

► R<sup>4</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl and R<sup>21</sup>;

► R<sup>5</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, fucosyl and R<sup>22</sup>;

► R<sup>7</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, arabinosyl and R<sup>23</sup>;

► R<sup>8</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub> and R<sup>24</sup>;

► R<sup>9</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, mannose, glycerol and R<sup>25</sup>;

► R<sup>10</sup>, R<sup>11</sup>, R<sup>17</sup> and R<sup>18</sup> represent, independently of each other, a substituent chosen from C<sub>1-6</sub>-alkyl and F;

► R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>24</sup> and R<sup>25</sup> represent, independently of each other, a substituent chosen from C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

Preferably, the compounds of formula (I) have one or other of the following characteristics, taken separately or in combination:

► n represents 2 or 3;

► A represents -C(O)- or -CH<sub>2</sub>-;

► B represents a phenylene;

► C represents -O-;

► D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;

► E and G represent NHC(O)CH<sub>3</sub>;

► R<sup>1</sup> represents H, CH<sub>3</sub> or C(O)CH<sub>3</sub>;

► R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;

► R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;

► R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl.

Among these compounds, the ones that are preferred are those of formula (I) simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH<sub>2</sub>-;
- ▶ E and G represent NHC(O)CH<sub>3</sub>;
- ▶ R<sup>1</sup> represents H, CH<sub>3</sub> or C(O)CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl;

even more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH<sub>2</sub>-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH<sub>3</sub>;
- ▶ R<sup>1</sup> represents H, CH<sub>3</sub> or C(O)CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl;

and most preferably the compounds of formula (I) simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH<sub>2</sub>-;
- ▶ C represents -O-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH<sub>3</sub>;
- ▶ R<sup>1</sup> represents H, CH<sub>3</sub> or C(O)CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl.

Among these preferred compounds, mention may be made of the compounds of formula (I) simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents -C(O)- or -CH<sub>2</sub>-;
- ▶ B represents a phenylene;

- C represents -O-;
- D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- E and G represent  $\text{NHC(O)CH}_3$ ;
- $\text{R}^1$  represents H,  $\text{CH}_3$  or  $\text{C(O)CH}_3$ ;
- $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$  and  $\text{R}^9$  represent H;
- $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl.

Among the compounds of the present invention, the compounds for which A represents a carbonyl group are particularly advantageous, and may be represented by formula (Ia):



in which

- n represents 1, 2 or 3,
- B represents
  - an arylene;
  - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a naphthylene;
  - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
  - a divalent radical derived from 2 fused heteroaromatic rings containing 5 or 6 atoms each, and comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a biphenylene;
  - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $\text{R}^{12}$  and  $\text{R}^{13}$  chosen, independently of each other, from halogen, CN,  $\text{C(O)OR}^{14}$ ,  $\text{C(O)NR}^{15}\text{R}^{16}$ ,  $\text{CF}_3$ ,  $\text{OCF}_3$ ,  $-\text{NO}_2$ ,  $\text{N}_3$ ,  $\text{OR}^{14}$ ,  $\text{SR}^{14}$ ,  $\text{NR}^{15}\text{R}^{16}$  and  $\text{C}_{1-6}\text{-alkyl}$ ;

- C represents a substituent chosen from -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>17</sup>-, -CR<sup>17</sup>R<sup>18</sup>-, -NH- and -NR<sup>19</sup>;
- D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;
- E and G represent, independently of each other, a substituent chosen from H, OH, OR<sup>20</sup>, NH<sub>2</sub> and NHR<sup>20</sup>;
- R<sup>1</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, C(O)H and C(O)CH<sub>3</sub>;
- R<sup>2</sup>, R<sup>3</sup> and R<sup>6</sup> represent, independently of each other, a substituent chosen from H, C<sub>1-6</sub>-alkyl, C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;
- R<sup>4</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl and R<sup>21</sup>;
- R<sup>5</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, fucosyl and R<sup>22</sup>;
- R<sup>7</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, arabinosyl and R<sup>23</sup>;
- R<sup>8</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub> and R<sup>24</sup>;
- R<sup>9</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, mannose, glycerol and R<sup>25</sup>;
- R<sup>10</sup>, R<sup>11</sup>, R<sup>17</sup> and R<sup>18</sup> represent, independently of each other, a substituent chosen from C<sub>1-6</sub>-alkyl and F;
- R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup> and R<sup>19</sup> represent, independently of each other, a substituent chosen from H, C<sub>1-6</sub>-alkyl, -C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;
- R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>24</sup> and R<sup>25</sup> represent, independently of each other, a substituent chosen from C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium, or tetraalkylammonium salts.

Among these compounds of formula (Ia) the ones that are preferred are those having the following characteristics, taken separately or in combination:

- n represents 2 or 3;
- B represents a phenylene;
- C represents -O-;
- D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- E and G represent NHC(O)CH<sub>3</sub>;

- ▶ R<sup>1</sup> represents H or CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl;

5 more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent NHC(O)CH<sub>3</sub>;
- ▶ R<sup>1</sup> represents H or CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl;

even more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH<sub>3</sub>;
- ▶ R<sup>1</sup> represents H or CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl.

Among these compounds of formula (Ia), the ones that are more preferred are those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ C represents -O-;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent NHC(O)CH<sub>3</sub>;
- ▶ R<sup>1</sup> represents H or CH<sub>3</sub>;
- ▶ R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>9</sup> represent H;
- ▶ R<sup>4</sup> represents H, C(O)CH<sub>3</sub> or C(O)NH<sub>2</sub>;
- ▶ R<sup>8</sup> represents H, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub>, fucosyl or methylfucosyl;


or those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ C represents -O-;



- D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- E and G represent  $\text{NHC(O)CH}_3$ ;
- $\text{R}^1$  represents H or  $\text{CH}_3$ ;
- $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$  and  $\text{R}^9$  represent H;
- $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl.

Among the compounds of the present invention, the compounds for which A represents a methylene group are also particularly advantageous, and may be represented by formula (Ib):



in which

- n represents 1, 2 or 3;
- B represents
  - an arylene;
  - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a naphthylene;
  - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a divalent radical derived from 2 fused aromatic rings each containing 5 or 6 atoms;
  - a divalent radical derived from 2 fused aromatic or heteroaromatic rings each containing 5 or 6 atoms, comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a biphenylene;
  - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $\text{R}^{12}$  and  $\text{R}^{13}$  chosen, independently of each other, from halogen, CN,  $\text{C(O)OR}^{14}$ ,  $\text{C(O)NR}^{15}\text{R}^{16}$ ,  $\text{CF}_3$ ,  $\text{OCF}_3$ ,  $-\text{NO}_2$ ,  $\text{N}_3$ ,  $\text{OR}^{14}$ ,  $\text{SR}^{14}$ ,  $\text{NR}^{15}\text{R}^{16}$  and  $\text{C}_{1-6}\text{-alkyl}$ ;

- C represents a substituent chosen from -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>17</sup>-, -CR<sup>17</sup>R<sup>18</sup>-, -NH- and -NR<sup>19</sup>;
- D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms;
- E and G represent, independently of each other, a substituent chosen from H, OH, OR<sup>20</sup>, NH<sub>2</sub> and NHR<sup>20</sup>;
- R<sup>1</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, C(O)H and C(O)CH<sub>3</sub>;
- R<sup>2</sup>, R<sup>3</sup> and R<sup>6</sup> represent, independently of each other, a substituent chosen from H, C<sub>1-6</sub>-alkyl, C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;
- R<sup>4</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl and R<sup>21</sup>;
- R<sup>5</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, fucosyl and R<sup>22</sup>;
- R<sup>7</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, arabinosyl and R<sup>23</sup>;
- R<sup>8</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl, SO<sub>3</sub>H, SO<sub>3</sub>Li, SO<sub>3</sub>Na, SO<sub>3</sub>K, SO<sub>3</sub>N(C<sub>1-8</sub>alkyl)<sub>4</sub> and R<sup>24</sup>;
- R<sup>9</sup> represents a substituent chosen from H, C<sub>1-6</sub>-alkyl, mannose, glycerol and R<sup>25</sup>;
- R<sup>10</sup>, R<sup>11</sup>, R<sup>17</sup> and R<sup>18</sup> represent, independently of each other, a substituent chosen from C<sub>1-6</sub>-alkyl and F;
- R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup> and R<sup>19</sup> represent, independently of each other, a substituent chosen from H, C<sub>1-6</sub>-alkyl, -C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;
- R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>24</sup> and R<sup>25</sup> represent, independently of each other, a substituent chosen from C(O)C<sub>1-6</sub>-alkyl, -C(S)C<sub>1-6</sub>-alkyl, -C(O)OC<sub>1-6</sub>-alkyl, -C(O)NH<sub>2</sub>, -C(S)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)NHC<sub>1-6</sub>-alkyl, -C(S)NHC<sub>1-6</sub>-alkyl and -C(NH)NHC<sub>1-6</sub>-alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

Among these compounds of formula (Ib) the ones that are preferred are those having the following characteristics, taken separately or in combination:

- n represents 2 or 3;
- B represents a phenylene;
- C represents -O-;
- D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- E and G represent NHC(O)CH<sub>3</sub>;

- ▶  $R^1$  represents H or  $C(O)CH_3$ ;
- ▶  $R^2, R^3, R^5, R^6, R^7$  and  $R^9$  represent H;
- ▶  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;
- ▶  $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

5 more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent  $NHC(O)CH_3$ ;
- ▶  $R^1$  represents H or  $C(O)CH_3$ ;
- ▶  $R^2, R^3, R^5, R^6, R^7$  and  $R^9$  represent H;
- 10 ▶  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;
- ▶  $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

even more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17
- 15 carbon atoms;
- ▶ E and G represent  $NHC(O)CH_3$ ;
- ▶  $R^1$  represents H or  $C(O)CH_3$ ;
- ▶  $R^2, R^3, R^5, R^6, R^7$  and  $R^9$  represent H;
- ▶  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;
- 20 ▶  $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl.

Among these compounds of formula (Ib), the ones that are more preferred are those simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ C represents  $-O-$ ;
- 25 ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent  $NHC(O)CH_3$ ;
- ▶  $R^1$  represents H or  $C(O)CH_3$ ;
- ▶  $R^2, R^3, R^5, R^6, R^7$  and  $R^9$  represent H;
- 30 ▶  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;
- ▶  $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

or those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- 35 ▶ C represents  $-O-$ ;

- D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- E and G represent  $\text{NHC(O)CH}_3$ ;
- $\text{R}^1$  represents H or  $\text{C(O)CH}_3$ ;
- $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$  and  $\text{R}^9$  represent H;
- $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-6}\text{alkyl})_4$ , fucosyl or methylfucosyl.

Among the compounds of the present invention, the compounds for which C represents an oxygen atom are also particularly advantageous, and may be represented by formula (Ic):



(Ic)

in which

- n represents 1, 2 or 3, preferably 2 or 3;
- A represents a substituent chosen from  $-\text{C(O)}-$ ,  $-\text{C(S)}-$ ,  $-\text{CH}_2-$ ,  $-\text{CHR}^{10}-$ ,  $-\text{CR}^{10}\text{R}^{11}-$ ,  $-\text{C(O)O}-$ ,  $-\text{C(O)S}-$ ,  $-\text{C(S)O}-$ ,  $-\text{C(S)S}-$ ,  $-\text{C(O)NH}-$ ,  $-\text{C(NH)NH}-$  and  $-\text{C(S)NH}-$ , preferably  $-\text{C(O)}-$ ;
- B represents
  - an arylene;
  - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a naphthylene;
  - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
  - a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each, comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a biphenylene;
  - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $C(O)OR^{14}$ ,  $C(O)NR^{15}R^{16}$ ,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl;

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms, preferably a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;

► E and G represent, independently of each other, a substituent chosen from H, OH,  $OR^{20}$ ,  $NH_2$  and  $NHR^{20}$ , preferably  $NHC(O)CH_3$ ;

►  $R^1$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl,  $C(O)H$  and  $C(O)CH_3$ , preferably H or  $CH_3$ ;

►  $R^2$ ,  $R^3$  and  $R^6$  represent, independently of each other, a substituent chosen from H,  $C_{1-6}$ -alkyl,  $-C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl; preferably H;

►  $R^4$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl and  $R^{21}$ , preferably H,  $C(O)CH_3$  or  $C(O)NH_2$ ;

►  $R^5$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, fucosyl and  $R^{22}$ , preferably H;

►  $R^7$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, arabinosyl and  $R^{23}$ , preferably H;

►  $R^8$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$  and  $R^{24}$ , preferably H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

►  $R^9$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, mannose, glycerol and  $R^{25}$ , preferably H;

►  $R^{10}$ ,  $R^{11}$ ,  $R^{17}$  and  $R^{18}$  represent, independently of each other, a substituent chosen from  $C_{1-6}$ -alkyl and F;

►  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$  and  $R^{19}$  represent, independently of each other, a substituent chosen from H,  $C_{1-6}$ -alkyl,  $-C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl;

►  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$  and  $R^{25}$  represent, independently of each other, a substituent chosen from  $C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones, metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

Among the compounds of formula (Ic), the ones that are preferred are those having one or other of the following characteristics, taken separately or in combination:

► n represents 2 or 3;

► A represents  $-C(O)-$  or  $-CH_2-$ ;

- ▶ B represents a phenylene;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H,  $\text{CH}_3$  or  $\text{C(O)CH}_3$ ;
- ▶  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- ▶  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl;

more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents  $-\text{C(O)}-$  or  $-\text{CH}_2-$ ;
- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H,  $\text{CH}_3$  or  $\text{C(O)CH}_3$ ;
- ▶  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- ▶  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl;

even more preferably, those simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ A represents  $-\text{C(O)}-$  or  $-\text{CH}_2-$ ;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H,  $\text{CH}_3$  or  $\text{C(O)CH}_3$ ;
- ▶  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- ▶  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl;

most preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ A represents  $-\text{C(O)}-$  or  $-\text{CH}_2-$ ;
- ▶ B represents a phenylene;
- ▶ D represents a linear hydrocarbon-based chain containing 11 carbon atoms, which is saturated, or unsaturated between carbons 4 and 5;
- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H,  $\text{CH}_3$  or  $\text{C(O)CH}_3$ ;
- ▶  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;

- $R^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl.

Among the compounds of the present invention, the compounds for which A represents a carbonyl group and C represents an oxygen atom are most particularly advantageous, and may be represented by formula (Id):



(Id)

in which

- $n$  represents 1, 2 or 3, preferably 2 or 3;
- B represents
- an arylene;
  - a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a naphthylene;
  - a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a divalent radical derived from 2 fused aromatic rings each containing 5 or 6 atoms;
  - a divalent radical derived from 2 fused aromatic or heteroaromatic rings each containing 5 or 6 atoms, comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
  - a biphenylene;
  - or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $\text{C}(\text{O})\text{OR}^{14}$ ,  $\text{C}(\text{O})\text{NR}^{15}\text{R}^{16}$ ,  $\text{CF}_3$ ,  $\text{OCF}_3$ ,  $-\text{NO}_2$ ,  $\text{N}_3$ ,  $\text{OR}^{14}$ ,  $\text{SR}^{14}$ ,  $\text{NR}^{15}\text{R}^{16}$  and  $\text{C}_{1-6}\text{-alkyl}$ ;

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms, preferably a linear hydrocarbon-based chain containing 11 carbon atoms, which is saturated, or unsaturated between carbons 4 and 5;

► E and G represent, independently of each other, a substituent chosen from H, OH,  $\text{OR}^{20}$ ,  $\text{NH}_2$  and  $\text{NHR}^{20}$ , preferably  $\text{NHC}(\text{O})\text{CH}_3$ ;

►  $R^1$  represents a substituent chosen from H,  $\text{C}_{1-6}\text{-alkyl}$ ,  $\text{C}(\text{O})\text{H}$  and  $\text{C}(\text{O})\text{CH}_3$ , preferably H or  $\text{CH}_3$ ;

►  $R^2$ ,  $R^3$  and  $R^6$  represent, independently of each other, a substituent chosen from H,  $C_{1-6}$ -alkyl,  $-C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl; preferably H;

►  $R^4$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl and  $R^{21}$ , preferably H,  $C(O)CH_3$  or  $C(O)NH_2$ ;

►  $R^5$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, fucosyl and  $R^{22}$ , preferably H;

►  $R^7$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, arabinosyl and  $R^{23}$ , preferably H;

►  $R^8$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$  and  $R^{24}$ , preferably H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

►  $R^9$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, mannose, glycerol and  $R^{25}$ , preferably H;

►  $R^{10}$ ,  $R^{11}$ ,  $R^{17}$  and  $R^{18}$  represent, independently of each other, a substituent chosen from  $C_{1-6}$ -alkyl and F;

►  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$  and  $R^{19}$  represent, independently of each other, a substituent chosen from H,  $C_{1-6}$ -alkyl,  $-C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl;

►  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$  and  $R^{25}$  represent, independently of each other, a substituent chosen from  $-C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones and metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

Among the compounds of formula (Id), the ones that are preferred are those having one or other of the following characteristics, taken separately or in combination;

► n represents 2 or 3;

► B represents a phenylene;

► D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;

► E and G represent  $NHC(O)CH_3$ ;

►  $R^1$  represents H or  $CH_3$ ;

►  $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^9$  represent H;

►  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;

►  $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

more preferably, those simultaneously having the following characteristics:

► n represents 2 or 3;



- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H or  $\text{CH}_3$ ;
- ▶  $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- ▶  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl;

even more preferably, those simultaneously having the following characteristics:


- ▶ n represents 2 or 3;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;

- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H or  $\text{CH}_3$ ;
- ▶  $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- ▶  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl;

and most preferably those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ D represents a linear hydrocarbon-based chain containing 11 carbon, which is saturated, or unsaturated between carbons 4 and 5;
- ▶ E and G represent  $\text{NHC(O)CH}_3$ ;
- ▶  $\text{R}^1$  represents H or  $\text{CH}_3$ ;
- ▶  $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6, \text{R}^7$  and  $\text{R}^9$  represent H;
- ▶  $\text{R}^4$  represents H,  $\text{C(O)CH}_3$  or  $\text{C(O)NH}_2$ ;
- ▶  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , fucosyl or methylfucosyl.

Among the compounds of the present invention, the compounds for which A represents a methylene group and C represents an oxygen atom are also most particularly advantageous, and may be represented by formula (Ie):



(Ie)

in which

► n represents 1, 2 or 3, preferably 2 or 3;

► B represents

- an arylene;
- a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a naphthylene;
- a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a divalent radical derived from 2 fused aromatic rings containing 5 or 6 atoms each;
- a divalent radical derived from 2 fused aromatic or heteroaromatic rings containing 5 or 6 atoms each, comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a biphenylene;
- or a heterobiphenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $C(O)OR^{14}$ ,  $C(O)NR^{15}R^{16}$ ,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl;

► D represents a linear or branched, saturated or unsaturated hydrocarbon-based chain containing from 2 to 20 carbon atoms, preferably a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;

► E and G represent, independently of each other, a substituent chosen from H, OH,  $OR^{20}$ ,  $NH_2$  and  $NHR^{20}$ , preferably  $NHC(O)CH_3$ ;

►  $R^1$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl,  $C(O)H$  and  $C(O)CH_3$ , preferably H or  $CH_3$ ;

►  $R^2$ ,  $R^3$  and  $R^6$  represent, independently of each other, a substituent chosen from H,  $C_{1-6}$ -alkyl,  $C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl; preferably H;

►  $R^4$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl and  $R^{21}$ , preferably H,  $C(O)CH_3$  or  $C(O)NH_2$ ;

- ▶  $R^5$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, fucosyl and  $R^{22}$ , preferably H;
- ▶  $R^7$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, arabinosyl and  $R^{23}$ , preferably H;
- ▶  $R^8$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$  and  $R^{24}$ , preferably H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;
- ▶  $R^9$  represents a substituent chosen from H,  $C_{1-6}$ -alkyl, mannose, glycerol and  $R^{25}$ , preferably H;
- ▶  $R^{10}$ ,  $R^{11}$ ,  $R^{17}$  and  $R^{18}$  represent, independently of each other, a substituent chosen from  $C_{1-6}$ -alkyl and F;
- ▶  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$  and  $R^{19}$  represent, independently of each other, a substituent chosen from H,  $C_{1-6}$ -alkyl,  $-C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl;
- ▶  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$  and  $R^{25}$  represent, independently of each other, a substituent chosen from  $C(O)C_{1-6}$ -alkyl,  $-C(S)C_{1-6}$ -alkyl,  $-C(O)OC_{1-6}$ -alkyl,  $-C(O)NH_2$ ,  $-C(S)NH_2$ ,  $-C(NH)NH_2$ ,  $-C(O)NHC_{1-6}$ -alkyl,  $-C(S)NHC_{1-6}$ -alkyl and  $-C(NH)NHC_{1-6}$ -alkyl;

and also the possible geometrical and/or optical isomers, enantiomers and/or diastereoisomers, tautomers, salts, N-oxides, sulfoxides, sulfones and metal or metalloid complexes thereof, which are agriculturally acceptable. Among the compounds defined above, the most important compounds are the salts, more particularly the lithium, sodium, potassium or tetraalkylammonium salts.

Among the compounds of (Ie), the ones that are preferred are those having one or other of the following characteristics, taken separately or in combination:

- ▶ n represents 2 or 3;
- ▶ B represents a phenylene;
- ▶ D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- ▶ E and G represent  $NHC(O)CH_3$ ;
- ▶  $R^1$  represents H or  $C(O)CH_3$ ;
- ▶  $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^9$  represent H;
- ▶  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;
- ▶  $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

more preferably, those simultaneously having the following characteristics:

- ▶ n represents 2 or 3;
- ▶ E and G represent  $NHC(O)CH_3$ ;
- ▶  $R^1$  represents H or  $C(O)CH_3$ ;
- ▶  $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^9$  represent H;
- ▶  $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;

- $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

even more preferably, those simultaneously having the following characteristics:

- n represents 2 or 3;
- D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 3 to 17 carbon atoms;
- E and G represent  $NHC(O)CH_3$ ;
- $R^1$  represents H or  $C(O)CH_3$ ;
- $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^9$  represent H;
- $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;

- $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl;

and most preferably those simultaneously having the following characteristics:

- n represents 2 or 3;
- B represents a phenylene;
- D represents a linear hydrocarbon-based chain containing 11 carbons, which is saturated, or unsaturated between carbons 4 and 5;
- E and G represent  $NHC(O)CH_3$ ;
- $R^1$  represents H or  $C(O)CH_3$ ;
- $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^9$  represent H;
- $R^4$  represents H,  $C(O)CH_3$  or  $C(O)NH_2$ ;
- $R^8$  represents H,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$ ,  $SO_3N(C_{1-8}alkyl)_4$ , fucosyl or methylfucosyl.

Among the compounds of formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) according to the invention, the ones that are preferred are those for which:

- B represents a substituent chosen from:

B1		B6		B11		B16	
B2		B7		B12		B17	
B3		B8		B13		B18	
B4		B9		B14		B19	
B5		B10		B15		B20	

in which  $R^{12}$  and  $R^{13}$  represent two substituents chosen, independently of each other, from halogen, CN,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl.

5 Among the compounds of formula (I), (Ia), (Ib), (Ic), (Id) or (Ie) according to the invention, the ones that are also preferred are those for which

► B represents

- an arylene;
- a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;
- a naphthylene;
- or a heteronaphthylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $C(O)OR^{14}$ ,  $C(O)NR^{15}R^{16}$ ,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl;

preferably, those for which

► B represents

- an arylene;
- or a heteroarylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $C(O)OR^{14}$ ,  $C(O)NR^{15}R^{16}$ ,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl;

more preferably, those for which

► B represents

- a phenylene;
- or a heterophenylene comprising 1 or 2 hetero atoms chosen from nitrogen, oxygen and sulfur;

these groups possibly being substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $C(O)OR^{14}$ ,  $C(O)NR^{15}R^{16}$ ,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl;

mention may be made especially of those for which

- B represents a phenylene B1 that may be substituted with one or two substituents  $R^{12}$  and  $R^{13}$  chosen, independently of each other, from halogen, CN,  $CF_3$ ,  $OCF_3$ ,  $-NO_2$ ,  $N_3$ ,  $OR^{14}$ ,  $SR^{14}$ ,  $NR^{15}R^{16}$  and  $C_{1-6}$ -alkyl.

Among the preferred compounds of the present invention, mention may also be made of those having one of the following characteristics, taken separately or in combination:

- $n = 2$  or  $3$ ;
- A represents  $-C(O)-$  or  $-CH_2-$ ;
- C represents  $-O-$ ;
- E and G represent  $NHC(O)CH_3$ ;
- $R^1$  represents H or  $C(O)CH_3$ ;
- $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$  and  $R^7$  represent a hydrogen atom;
- $R^4$  represents a substituent chosen from H,  $C(O)CH_3$  and  $C(O)NH_2$ ;
- $R^8$  represents a substituent chosen from H, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl,  $SO_3H$ ,  $SO_3Li$ ,  $SO_3Na$ ,  $SO_3K$  and  $SO_3N(C_{1-8}alkyl)_4$ ;
- $R^9$  represents a hydrogen atom;

even more preferably, those having the following combination of characteristics:

- $n = 2$  or  $3$ ;
- A represents  $-C(O)-$  or  $-CH_2-$ ;
- C represents  $-O-$ ;

► D represents a linear, saturated or unsaturated hydrocarbon-based chain containing from 7 to 15 carbon atoms; preferably a hydrocarbon-based chain according to one of the formulae represented below

D1		D4	
D2		D5	
D3		D6	

5 in which

- $m = 1$  to 12
- $p = 0$  to 11
- $q = 6$  to 14
- $s = 5$  to 13

10 with  $m+p \leq 12$  and  $m+p \geq 4$ ; even more preferably a hydrocarbon-based chain according to one of the formulae represented below


D1	
D2	
D3	

in which

- 15
- $m = 1$  to 12
  - $p = 0$  to 11
  - $q = 6$  to 14
  - with  $m+p \leq 12$  and  $m+p \geq 4$ ; and most preferably a linear hydrocarbon-based chain containing 11 carbon atoms, which is saturated, or unsaturated between carbon atoms 4 and 5;

- E and G represent  $\text{NHC(O)CH}_3$ ;
- $\text{R}^1$  represents H or  $\text{C(O)CH}_3$ ;
- $\text{R}^2, \text{R}^3, \text{R}^5, \text{R}^6$  and  $\text{R}^7$  represent a hydrogen atom;
- $\text{R}^4$  represents a substituent chosen from H,  $\text{C(O)CH}_3$  and  $\text{C(O)NH}_2$ ;
- $\text{R}^8$  represents a substituent chosen from H, fucosyl, methylfucosyl, sulfofucosyl, acetylfucosyl, arabinosyl,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$  and  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ ;
- $\text{R}^9$  represents a hydrogen atom;


in particular, the compounds for which  $\text{R}^8$  represents H,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$ ,  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$  or a substituent of formula:







in which





- $\text{R}^{26}$  represents a substituent chosen from H and  $\text{CH}_3$ , preferably H;
- $\text{R}^{27}$  and  $\text{R}^{28}$  represent, independently of each other, a substituent chosen from H,  $\text{C(O)CH}_3$ ,  $\text{SO}_3\text{H}$ ,  $\text{SO}_3\text{Li}$ ,  $\text{SO}_3\text{Na}$ ,  $\text{SO}_3\text{K}$  and  $\text{SO}_3\text{N}(\text{C}_{1-8}\text{alkyl})_4$ , preferably  $\text{R}^{27}$  and  $\text{R}^{28}$  represent H.

As examples of compounds according to the invention that are particularly advantageous and preferred, mention may be made of the compounds corresponding to one of the following formulae:









in which, when it is present, M represents a cation chosen from  $H^+$ ,  $Li^+$ ,  $Na^+$ ,  $K^+$  and  $(C_{1-8}alkyl)_4N^+$ .


Besides the compounds of the invention that have just been specifically described, the variants of combinations of possible substituents for the formulae (I), (Ia), (Ib), (Ic), (Id) and (Ie) especially, also form part of the invention.

It is known that a chitin oligomer not containing a lipid chain is not active, and that the degradation of the Nod factors by breaking the amide bond in the rhizosphere thus leads to a loss of activity.


In order to limit, or even prevent, this degradation, a series of analogous compounds, some of which are more stable than the natural Nod factors, was prepared.

## II-1. STRUCTURE OF COMPOUNDS ACCORDING TO THE INVENTION

Compounds containing a meta-substituted benzamide group were prepared. It is preferred to keep identical the total number of atoms along the chain (16) and also the unsaturation of cis type in position 9. In practice, for the production of the starting materials, the lipid chain may be linked to the aromatic ring via an oxygen atom.





An analog 4 containing a meta-substituted benzylamine function, and also an N-acetylated analog 5, which makes it possible to regain the overall charge of the natural product, were also synthesized. These analogs were prepared in the sulfated series.





- 4  $R^1 = H$   
5  $R^1 = Ac$


Two other sulfated analogs, one containing a fully saturated chain and the other an unsaturated chain of alkyne type, make it possible to study the effect of the unsaturation of Z type in position 9 present on the natural product.



Finally, two sulfated analogs, the substitution on the aromatic ring of which is in the ortho position for one and in the para position for the other, make it possible to study the effect of the unsaturation of trans type located in position 2 on the natural product.




Finally, an analog, derived from a fucosyl pentamer, bearing a meta substitution on the chain, was prepared.



**10**

The references for the biological tests are the following compounds:




## II-2. SYNTHESIS OF THE VARIOUS AROMATIC CHAINS

For the benzamide LCOs, the coupling with the amino tetramer is performed with a benzoyl chloride (acylation) and for the benzyl LCOs, with a benzaldehyde (reductive alkylation).


### II-2.1. Synthesis of aromatic chains meta-substituted with the undec-4Z-enoxy chain

According to the reaction scheme below, the methyl ester **15** is prepared, from which reduction to the aldehyde or saponification to the acid (acyl chloride precursor) may be envisaged.


To do this, 1-iodoundec-4Z-ene **13** is used to alkylate methyl 3-hydroxybenzoate. The ester **15** is isolated in a yield of 76% .



Conversion of the ester to the aldehyde **17** is performed in two steps.





Moreover, the acyl chloride **19** is obtained by saponification of the ester **15** followed by reaction with oxalyl chloride.



## II-2.2. Synthesis of aromatic chains meta-substituted with undecanyloxy and undec-4-ynyloxy chains


- 5 The same procedure with 1-bromoundecane or 1-iodoundec-4-yne in anhydrous DMF, followed by saponification and formation of the chloride, lead to the acid chlorides **23** and **27**.






### II-2.3. Synthesis of aromatic chains ortho- or para-substituted with the undec-4Z-enyloxy chain

The acid chlorides **31** and **35** are similarly prepared from **29** and **33**, which are obtained as previously by Williamson coupling of 1-iodoundec-4Z-ene **13** with methyl 2-hydroxybenzoate **28** (or methyl salicylate) in a yield of 66%, and with methyl 4-hydroxybenzoate **32** in a yield of 79%.









The saponifications and conversions to chloride are quantitative in both cases.

5 II-3. N-acylation of the sulfated tetramer **CO-IV(NH<sub>2</sub>, S)** with the various benzoyl chlorides



10 II-3.1. Coupling with 3-(undec-4Z-enyloxy)benzoyl chloride **19**




15

The coupling may be performed by dissolving the starting material in a DMF-water mixture in the presence of sodium hydrogen carbonate. Under these conditions, only the free amine is acylated. With





For these two analogs, by adopting a similar protocol, a yield of 48% is obtained for the ortho-substituted derivative **8** and a yield of 40% is obtained for the para-substituted derivative **9**. For the two reactions, 4 equivalents of chloride were used. The purity is also checked by HPLC.

5



II-4. N-Acylation of the nonsulfated tetramer **CO-IV(NH<sub>2</sub>)** with 3-(undec-4Z-enyloxy)benzoyl chloride **19**

10





15

The reaction was carried out as previously in a DMF-water mixture, in which the starting material and the chloride are soluble. In order to facilitate the final purification, the reaction is performed in the presence of a basic Dowex resin ( $\text{HCO}_3^-$ ).

5 At the end of the reaction, the reaction medium is diluted with an acetonitrile/water mixture, and the expected compound is purified by filtration of the resin, passing through acidic Dowex ( $\text{H}^+$ ), resin, concentration and washing of the solid residue with ethyl acetate and then with water. 22% of the expected product **2** are thus isolated.

10 II-5. N-acylation of the fucosylated pentamer **CO-V(NH<sub>2</sub>, Fuc)** with 3-(undec-4Z-enyloxy)benzoyl chloride **19**




**10**

15 The reaction was performed as for the preceding product, in a DMF-water mixture, in which the starting material and the chloride are soluble. In order to facilitate final purification, the reaction is performed in the presence of a basic Dowex resin ( $\text{HCO}_3^-$ ).

20 At the end of the reaction, the reaction medium is diluted with an acetonitrile/water mixture, and the expected compound is purified by filtration of the resin, passage through acidic Dowex resin ( $\text{H}^+$ ), concentration and washing of the solid residue with ethyl acetate and then with water. 28% of the expected product **10** are thus isolated.


## II-6. Reductive alkylation of the sulfated tetramer with 3-(undec-4Z-enyloxy)benzaldehyde

### II-6.1. Alkylation of the tetramer CO-IV(NH<sub>2</sub>, S)



The reductive alkylation reaction was performed in anhydrous DMF in the presence of lithium bromide. With 12 equivalents of aldehyde and 15 equivalents of sodium cyanoborohydride, 71% of expected coupling product 4 are isolated by chromatography on silica gel after 24 hours.

### II-6.2. N-Acetylation of the coupling product obtained from the reductive alkylation



The reaction is performed in an ethyl acetate-methanol-water mixture by addition of acetic anhydride, in the presence of sodium hydrogen carbonate. After 12 hours, the starting material 4 is removed by passage through H<sup>+</sup> resin. After purification on silica, the expected product 5 is isolated in a yield of 77%. The purity is checked by HPLC.

## II-7. ACTIVITY TESTS

### II-7.1 Activity tests on temperate legumes of the Galegoid group

Temperate legumes of the Galeoid group are nodulated by rhizobia that produce Nod factors with the hydrophobic chain having a double bond conjugated to the carbonyl group. This group includes important legume crops such as alfalfa, pea, broad bean, chickpea and clover.

- 5 The sulfated products are tested on alfalfa for induction of the formation of root nodules, and on the model legume *Medicago truncatula* for induction of the expression of a symbiotic gene coding for an early nodulin.


## II-7.1. 1 Nodulation tests on alfalfa

10

Alfalfa plantlets are grown under axenic conditions in test tubes on a nitrogen-poor agar medium (Demont-Caulet et al., Plant Physiol., 120, 83-92, 1999). Untreated plantlets serve as control. Natural Nod or synthetic LCO factors are added at the concentrations indicated.

### 15 II-7.1.1.1 Results of nodulation tests


The benzamide derivative 3 meta-substituted with the undec-4Z-enyloxy chain shows advantageous activity, the activity being similar to that of the sulfated tetramer 11 acylated with the reference C16:1Δ9Z chain.



20


A: 11; B: 3; Control: no LCO

The benzyl derivative 4 has moderate activity relative to the benzamide derivative 3.



A: 11; B: 3; C: 4; Control: no LCO

Finally, the N-acetylation of the benzyl derivative 4 leads to an improvement in the response, but the activity remains lower than that of the benzamide derivative 3.




A: 3 to  $10^{-7}$ M; B: 4 to  $10^{-7}$ M; C: 5 to  $10^{-7}$ M; D: 11 to  $10^{-7}$ M;

E: 11 to  $10^{-9}$ M; Control: no LCO

- 5 These results indicate the importance of the amide bond. The results obtained with these benzamide derivatives confirm the effect of the amide-double bond conjugation present on the natural compound. The benzamide 7 meta-substituted with the undec-4-ynyloxy chain shows activity comparable to that of the benzamide derivative 3, whereas the benzamide compound 6 substituted with the fully saturated chain shows slightly lower activity. These results indicate that an unsaturation in position 4 may lead to an
- 10 increase in activity.





A: 3; B: 9; C: 8; D: 7; E: 6; Control: no LCO

The tests relating to the benzamide derivatives 8 and 9 ortho- and para-substituted with the undec-4Z-enyloxy chain reveal analogs that are less active than the meta-substituted benzamide derivative 3. The meta substitution is thus preferred as mimic for an unsaturation of trans type.

#### II-7.1.2. Tests of induction of early nodulin


These tests are performed to determine whether the synthetic LCOs induce symbiotic responses by activation of the same signal transduction pathway as the natural Nod factors. The tests are performed on the model legume *Medicago truncatula*. The activity of the sulfated benzamide derivative 3, meta-substituted with the undec-4Z-enyloxy chain, which is the most active synthetic compound in the nodulation test on alfalfa, is studied on "wild-type" plants and on a mutant in the gene *DMI1* which is altered in the transduction of the Nod factor signal (Catoira et al. *Plant Cell*, 12, 1647-1665, 2000). The compound that serves as reference is the sulfated tetramer 12 acylated with the C16:2Δ2E,9Z chain, which is an analog of the natural Nod factor. The control is the plant cultivated in the absence of LCO.

##### II-7.1.2.1 Reporter gene

It is generally difficult to determine the regulation of expression of a particular gene, during a biological process, since most of the specific products of these genes are not readily detectable or measurable. To overcome this problem, a technique of fusion with "reporter genes" is used, i.e. genes coding for a readily assayable protein. The fusion consists in combining the DNA sequence containing the gene regulatory regions that it is desired to study, with the DNA sequence of the reporter gene. The assembly is then reintroduced into the plant by transformation. Thus, if the target gene is expressed, the reporter gene is automatically expressed. It is then a matter of assaying the reporter gene protein.

In order to avoid a negative interaction with the activity of the plant, reporter genes that do not code for any enzyme normally formed by the plants are used. One of the enzymes most commonly used is  $\beta$ -glucuronidase (GUS) from *Escherichia coli*, a hydrolase that catalyzes the cleavage of a large variety of  $\beta$ -glucuronides. As commercial substrate of this enzyme, it is possible to use:

X-Gluc (Sigma B-4782): 5-bromo-4-chloro-3-indolyl glucuronide; the anion formed has a blue color.



#### II-7.1.2.2 Enod11::GUSA


The genes for the legumes involved in modulation may be classified into two major types: early nodulin genes (*ENOD*), which are activated in the first days of the infection and activation of the nodulation process; late nodulin genes, which are not activated until several days after the application of the bacteria, and do not intervene until the period of maturation of the nodules.

A new gene of *Medicago truncatula*, *MtENOD11*, coding for an RPRP (Repetitive proline-rich protein), and transcribed during the first steps of infection of nodulation on the nodule roots and tissues was identified (Journet et al. *Mol. Plant-Microbe Interact.*, 14, 737-748, 2001). Using the transgenic *Medicago truncatula* plant expressing the fusion *MtENOD11::GUSA*, it is possible to determine whether a Nod factor analog added to the culture medium of the plant has induced transcription of the *ENOD11* gene.

For the *ENOD11* transcription tests, a Fahraeus medium is used as for the modulation tests, but without agar. The seedlings are placed on paper in pockets containing the culture medium. The responses of two

types of transgenic plants bearing the *MtENOD11::GUS* fusion are compared: a "wild-type" (WT) Jemalong plant and a plant bearing a mutation in the *DMI1* gene, which is incapable of transducing the Nod factor signal. The plants are left to grow for 5 days and the plantlets are then treated with various concentrations of LCO. After 6 hours, the plantlets are removed and placed in aqueous medium containing X-Gluc for 1 to 2 hours. The number of roots giving a characteristic blue response is then counted.

This test is relatively sensitive, to the extent that it is possible to work at LCO concentrations that are lower than those for the nodulation tests.



A: 3; B: 12; Control: no LCO

It is found that the benzamide derivative 3 is approximately 10 times less active than the reference compound, the acylated tetramer 12. Moreover, as for the reference compound 12, the benzamide derivative 3 does not induce any response in the plants bearing the *DMI1* mutation. It


may thus be concluded that the synthetic compound of benzamide type activates the transcription of the ENOD11 gene via the same transduction pathway as that activated by the natural Nod factors.

5

### III EXAMPLES

For the aromatic derivatives, the ring is numbered according to the official nomenclature.


For the description of the NMR spectra for the CO and LCO, the sugars are numbered starting with the  
10 reducing end:



The conventional numbering is adopted on each sugar.

15

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(N-3-(undec-4Z-enyloxy)benzoyl)amino- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-D-glucopyranose (2)**



20

7.2 mg of **CO-IV(NH<sub>2</sub>)** are dissolved in 200  $\mu$ L of water and 500  $\mu$ L of DMF, and are then heated to 40°C. 36 mg of Dowex 1x2-100 resin (HCO<sub>3</sub><sup>-</sup>) are then added, followed by addition of 160  $\mu$ L of a solution of **19** in distilled THF (26  $\mu$ mol). 108 mg of HCO<sub>3</sub><sup>-</sup> resin and 480  $\mu$ L of the solution of **19** in distilled THF (78  $\mu$ mol) are added in three portions over 48 hours. The reaction medium is diluted with 3 mL of 1/1 acetonitrile/water mixture, the reaction medium is collected, leaving the resin, and is then filtered through  
25 cotton wool to remove the entrained resin beads. The filtrates are passed through a Dowex 50x8-100

resin (H<sup>+</sup>) and then concentrated, and washing of the solid residue is then performed with ethyl acetate, and then with water. 2 mg of a white powder are obtained, i.e. a yield of 22 %.

5 <sup>1</sup>H NMR (400 MHz, 20/1 DMSO-d<sub>6</sub>/D<sub>2</sub>O) δ (ppm):


7.40 – 7.31 (m, 3H, ArH-2, ArH-6 and ArH-5), 7.04 (m, 1H, ArH-4), 5.41-5.35 (m, 2H, CH=CH), 4.87 (d, 0.7H, J<sub>1,2</sub> = 2.3 Hz, H-1α<sup>I</sup>), 4.52 (d, 1H, J = 8.3 Hz, H-1β<sup>IV</sup>), 4.42 (d, 0.3H, J = 8.0 Hz, H-1β<sup>I</sup>), 4.33 (2d, 2H, J = 8.3Hz, H-1β<sup>II-III</sup>), 3.98 (t, 2H, J = 6.0 Hz, ArOCH<sub>2</sub>-CH<sub>2</sub>). 3.78 – 3.05 (m, 24H, other sugar Hs), 2.16 (dt, 2H, J = 5.8 and J = 6.7 Hz, CH<sub>2</sub>-CH=CH), 1.97 (dt, 2H, J = 6.0 and J = 6.2 Hz, CH=CH-CH<sub>2</sub>), 1.81 / 1.81/  
10 1.79 (3s, 9H, 3 COCH<sub>3</sub>), 1.80 – 1.72 (m, 2H, ArOCH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>), 1.28 – 1.13 (m, 8H, 4 CH<sub>2</sub>), 0.81 (t, 3H, CH<sub>3</sub>, J = 6.5 Hz).

Mass spectrum:

Positive electrospray (ESI) ionization m/z = 1183.5 [M + Na]<sup>+</sup>

15

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(N-3-(undec-4Z-  
enyloxy)benzoyl)amino-β-D-glucopyranosyl]-β-D-glucopyranosyl]-2-deoxy-β-D-glucopyranosyl}-2-  
deoxy-6-O-sulfo-D-glucopyranose, sodium salt (3)**



20

15 mg of **CO-IV(NH<sub>2</sub>,S)** (17 μmol) are dissolved in 100 μL of water and 250 μL of DMF. 3 mg of sodium hydrogen carbonate (34 μmol) are then added, followed by addition of 20 μL of a solution of **19** in THF at a concentration of 0.25 g/mL (16.4 μmol). The reaction medium is heated to 60°C and 100 μL of the solution of **48** and 10 mg of sodium hydrogen carbonate are added in six portions over 18 hours. After  
25 concentrating, the residue is purified by placing it in dichloromethane(DCM)/methanol (5/1) on a column of silica, while diluting it greatly, in order to remove the lipid chain. The elution is then performed with E/M/W (7/2/1 Ethyl acetate/Methanol/Water). 6.5 mg of a white solid are thus isolated, i.e. a yield of 33%.

<sup>1</sup>H NMR (400 MHz, DMSO-CD<sub>3</sub>OD (1/2)) δ (ppm):

7.48 and 7.41 (m, 2 H, ArH-2 and ArH-6), 7.36 (dd, 1 H, ArH-5,  $J_{5,6}$  7.7 Hz and  $J_{5,4}$  8.1 Hz), 7.07 (ddd, 1 H, ArH-4,  $J_{4,2} \approx J_{4,6}$  1.4 Hz), 5.41 (m, 2 H, CH=CH), 5.03 (d, 0.8 H, H-1 $\alpha$ <sup>I</sup>,  $J_{1\alpha,2}$  3.2 Hz), 4.68-4.59-4.50 (3 d, 3 H, H-1 $\beta$ <sup>II,III,IV</sup>,  $J_{1\beta,2}$  8.4 Hz, 8.5 Hz and 8.7 Hz), 4.56 (d, 0.2 H, H-1 $\beta$ <sup>I</sup>,  $J_{1\beta,2}$  7.7 Hz), 4.25-3.30 (m, 26 H, CH<sub>2</sub>-OAr, other Hs of the sugars), 2.25 (td, 2 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>,  $J$  6.7 Hz and  $J$  6.2 Hz), 2.10-1.90 (m, 11 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub> and 3 CH<sub>3</sub>CO), 1.83 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>,  $J$  6.7 Hz), 1.35-1.20 (m, 8 H, 4 CH<sub>2</sub>), 0.88 (m, 3 H, CH<sub>3</sub>)


Mass spectrum:

Negative ESI  $m/z$  = 1139.4 [M-Na]<sup>-</sup>

UV: 289 nm

Fluorescence:  $\lambda_{ex}$ : 289 nm;  $\lambda_{em}$ : 345 nm

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(*N*-3-(undec-4Z-  
enyloxy)benzyl)amino- $\beta$ -D-glucopyranosyl)- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-6-O-sulfo-D-glucopyranose, sodium salt (**4**)**



11 mg of **CO-IV(NH<sub>2</sub>,S)** (12  $\mu$ mol) are dissolved in 0.5 mL of DMF to which are added 12 mg of lithium bromide. 2 mg of sodium cyanoborohydride (32  $\mu$ mol) and 100  $\mu$ L of a solution of **17** in THF at a concentration of 73 mg/mL (26  $\mu$ mol) are added. The reaction medium is heated at 40°C for 4 hours. Every 2 hours, 2 equivalents of aldehyde and 2.5 equivalents of sodium cyanoborohydride are added, i.e. in total 12 equivalents of aldehyde and 15 equivalents of sodium cyanoborohydride. Although the conversion is not complete, the reaction is stopped by destroying the excess sodium cyanoborohydride with 0.5 N hydrochloric acid. When the evolution of gas has ended, the medium is diluted in water and freeze-dried. The resulting material is taken up in water, 5 mg of sodium hydrogen carbonate (59  $\mu$ mol) are added to return to basic pH, and the resulting material is then coevaporated twice with methanol. The residual white solid is placed in DCM/methanol (5/1) on a column of silica, while diluting it greatly, in order

to remove the lipid chain. The elution is then performed with E/M/W (5/2/1) and then (4/1/1). 10 mg of white needles are thus isolated, i.e. a yield of 71%.

$^1\text{H}$  NMR (400 MHz, DMSO- $\text{CD}_3\text{OD}$  (2/1))  $\delta$  (ppm):

7.31 (dd, 1 H, ArH-5,  $J_{4,5}$  8.2 Hz and  $J_{5,6}$  7.8 Hz), 7.02 (m, 2 H, ArH-2 and ArH-6), 6.90 (dd, 1 H, ArH-4,  $J_{4,6}$  2.3 Hz), 5.51 (m, 2 H, CH=CH), 5.08 (d, 0.8H, H-1 $\alpha^I$ ,  $J_{1\alpha,2}$  3.1 Hz), 4.67 (m, 2.2 H, H-1 $\beta^{I,II,III}$ ), 4.47 (d, 1 H, H-1 $\beta^{IV}$ ,  $J_{1\beta,2}$  8.0 Hz), 4.06 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 3.94 (s, 2 H, NH-CH<sub>2</sub>-Ar), 4.25 – 3.45 (m, 23 H, other Hs of the sugars), 2.45 (dd, 1 H, H<sub>2</sub><sup>IV</sup>,  $J_{1\beta,2} \approx J_{2,3}$  8.8 Hz), 2.31-2.12 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 2.07-2.04-2.01 (3 s, 9 H, 3 CH<sub>3</sub>CO), 1.89 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH=CH,  $J$  6.9 Hz), 1.45-1.25 (m, 8 H, 4 CH<sub>2</sub>), 0.97 (t, 3 H, CH<sub>3</sub>,  $J$  6.8 Hz)


$^{13}\text{C}$  NMR (50 MHz, DMSO- $\text{CD}_3\text{OD}$  (2/1))  $\delta$  (ppm):

172 (3 CH<sub>3</sub>CO), 160 (ArC-3), 132-131-130 (ArC-1, ArC-5, CH=CH), 122 (ArC-6), 115 (ArC-2, ArC-4), 105 (C-1 $\beta^{II,III,IV}$ ), 98 (C-1 $\beta^I$ ), 92 (C-1 $\alpha^I$ ), 82 – 53 (21 C of the sugars and Ar-CH<sub>2</sub>-NH), 68 (CH<sub>2</sub>-OAr), 33-23 (10 CH<sub>2</sub> and 3 CH<sub>3</sub>CO), 14 (CH<sub>3</sub>)

Mass spectrum:

Negative ESI  $m/z$  = 1125.4 [M-Na]<sup>-</sup>

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(N-3-(undec-4Z-enoxy)benzyl)acetamido- $\beta$ -D-glucopyranosyl)]- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-6-O-sulfo-D-glucopyranose, sodium salt (5)**



20 mg of sodium hydrogen carbonate and 15  $\mu\text{L}$  of acetic anhydride are added to a solution of 13 mg of **4** (11  $\mu\text{mol}$ ) in 0.3 mL of E/M/W (1/1/1). The reaction medium is stirred at room temperature for 12 hours. After concentrating, the residual oil is taken up in E/M/W (1/1/1) and Dowex 50x8-100 H<sup>+</sup> resin is added. The mixture is filtered and Amberlite IR120 Na<sup>+</sup> resin is added to the filtrate. After filtering and

concentrating, the product is purified by chromatography in E/M/W (4/1/1). 10 mg of a white solid are thus isolated, i.e. a yield of 77%.


$^1\text{H}$  NMR (400 MHz, DMSO- $\text{CD}_3\text{OD}$  (2/1))  $\delta$  (ppm):

7.25-7.18 (2 t, 1 H, ArH-5,  $J_{5,4}$  7.8 Hz and  $J_{5,6}$  7.9 Hz), 7.10-6.85 (m, 2 H, ArH-2 and ArH-6), 6.82-6.75 (2 d, 1 H, ArH-4), 5.40 (m, 2 H, CH=CH), 5.06 (d, 0.6 H, H-1 $\alpha$ <sup>I</sup>,  $J_{1\alpha,2}$  3.4 Hz), 4.75-4.35 (m, 3.4 H, H-1 $\beta$ <sup>I,II,III,IV</sup>), 4.30-4.05 (m, 2 H, H-6a,b<sup>I</sup>), 4.00 – 3.30 (m, 25 H, other Hs of the sugars and CH<sub>2</sub>-OAr), 3.80 (s, 2 H, NAc-CH<sub>2</sub>-Ar), 2.90 (m, 1 H, H-2<sup>IV</sup>), 2.23-2.03 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.99-1.90 (m, 12 H, CH<sub>3</sub>CO), 1.80 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH=CH,  $J$  6.9 Hz), 1.35-1.20 (m, 8 H, 4 CH<sub>2</sub>), 0.87 (m, 3 H, CH<sub>3</sub>)

$^{13}\text{C}$  NMR (50 MHz, DMSO- $\text{CD}_3\text{OD}$  (2/1))  $\delta$  (ppm): 176 (CH<sub>3</sub>CON), 174-173-173 (3 CH<sub>3</sub>CO), 161 (ArC-3), 141 (ArC-1), 132-130-129-127 (ArC-2, ArC-4, ArC-5, ArC-6, CH=CH), 103 (3 C-1 $\beta$ <sup>I,II,IV</sup>), 100 (C-1 $\beta$ <sup>I</sup>), 92 (C-1 $\alpha$ <sup>I</sup>), 82 – 50 (24 C of the sugars, Ar-CH<sub>2</sub>-NH and CH<sub>2</sub>-OAr), 33-23 (10 CH<sub>2</sub> and 3 CH<sub>3</sub>CO), 14 (CH<sub>3</sub>)

Mass spectrum: Negative ESI  $m/z$  = 1067.4 [M-Na]<sup>-</sup>

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(N-3-(undecanyloxy)benzoyl)amino- $\beta$ -D-glucopyranosyl)-2-deoxy- $\beta$ -D-glucopyranosyl]- $\beta$ -D-glucopyranosyl)-2-deoxy-6-O-sulfo-D-glucopyranose, sodium salt (**6**)**



15 mg of **CO-IV(NH<sub>2</sub>,S)** (17  $\mu\text{mol}$ ) are dissolved in 100  $\mu\text{L}$  of water and 250  $\mu\text{L}$  of DMF. 6 mg of sodium hydrogen carbonate (71  $\mu\text{mol}$ ) and then 25  $\mu\text{L}$  of a solution of **23** in THF at a concentration of 210 mg/mL (17  $\mu\text{mol}$ ) are then added. The reaction medium is heated to 60°C and 200  $\mu\text{L}$  of the solution of chloride and 16 mg of sodium hydrogen carbonate are added in eight portions over 24 hours. After concentrating, the residue is purified by placing it in DCM/methanol (5/1) on a column of silica, while diluting it greatly, in order to remove the lipid chain. The elution is then performed with E/M/W (4/1/1). 6.3 mg of a white solid are thus isolated, i.e. a yield of 32%.




<sup>1</sup>H NMR (400 MHz, DMSO-CD<sub>3</sub>OD (1/3)) δ (ppm): 7.44 (m, 2 H, ArH-2 and ArH-6), 7.39 (dd, 1 H, ArH-5,  $J_{5,4} \approx J_{5,6}$  7.9 Hz), 7.10 (ddd, 1 H, ArH-4,  $J_{4,6} \approx J_{4,2}$  2.1 Hz), 5.05 (d, 0.7 H, H-1 $\alpha^I$ ,  $J_{1\alpha,2}$  3.0 Hz), 4.70-4.40 (m, 3.3 H, H-1 $\beta^{I,II,III,IV}$ ), 4.22 (m, 1 H, H-6a<sup>I</sup>), 4.10-3.20 (m, 24 H, CH<sub>2</sub>-OAr and other Hs of the sugar), 2.03-1.99-1.96 (3 s, 9 H, CH<sub>3</sub>CO), 1.80 (m, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>), 1.35-1.25 (m, 8 H, 4 CH<sub>2</sub>), 0.92 (t, 3 H, CH<sub>3</sub>,  $J$  6.5 Hz)

Mass spectrum: Negative ESI

$m/z$  = 1141.5 [M-Na]-

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(N-3-(undec-4Z-nyloxy)benzoyl)amino- $\beta$ -D-glucopyranosyl)]- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-6-O-sulfo-D-glucopyranose, sodium salt (**7**)**



14 mg of **CO-IV(NH<sub>2</sub>,S)** (16  $\mu$ mol) are dissolved in 100  $\mu$ L of water and 250  $\mu$ L of DMF. 5 mg of sodium hydrogen carbonate (60  $\mu$ mol) and then 25  $\mu$ L of a solution of **27** in THF at a concentration of 190 mg/mL (16  $\mu$ mol) are then added. The reaction medium is heated to 60°C and 200  $\mu$ L of the solution of chloride and 16 mg of sodium hydrogen carbonate are added in eight portions over 24 hours. After concentrating, the residue is purified by placing it in DCM/methanol (5/1) on a column of silica, while diluting it greatly, in order to remove the lipid chain. The elution is then performed with E/M/W (4/1/1). 5.7 mg of expected product are thus isolated in the form of a white solid, i.e. a yield of 31%.

<sup>1</sup>H NMR (400 MHz, DMSO-CD<sub>3</sub>OD (1/2)) δ (ppm): 7.43 (m, 2 H, ArH-2 and ArH-6), 7.37 (dd, 1 H, ArH-5,  $J_{5,4}$  8.1 Hz and  $J_{5,6}$  8.0 Hz), 7.10 (ddd, 1 H, ArH-4,  $J_{4,2} \approx J_{4,6}$  2.0 Hz), 5.04 (d, 0.7 H, H-1 $\alpha^I$ ,  $J_{1\alpha,2}$  3.3 Hz), 4.65-4.59 (2 d, 2 H, H-1 $\beta^{II,III}$ ,  $J_{1\beta,2}$  8.4 Hz and  $J_{1\beta,2}$  8.5 Hz), 4.54 (d, 0.3 H, H-1 $\beta^I$ ,  $J_{1\beta,2}$  7.9 Hz), 4.49 (d, 1 H, H-1 $\beta^{IV}$ ,  $J_{1\beta,2}$  8.7 Hz), 4.23 (dd, 1 H, H-6a<sup>I</sup>,  $J_{6a,6b}$  11.1 Hz and  $J_{6a,5}$  3.7 Hz), 4.12 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.2 Hz), 4.10-3.40 (m, 21 H, other Hs of the sugars), 2.35-2.13 (2 m, 4 H, CH<sub>2</sub>-C $\equiv$ C-CH<sub>2</sub>), 2.02-1.98-1.96 (3 s, 9 H, 3 CH<sub>3</sub>CO), 1.92 (m, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>), 1.45-1.25 (m, 8 H, 4 CH<sub>2</sub>), 0.88 (t, 3 H, CH<sub>3</sub>,  $J$  6.7 Hz)


<sup>13</sup>C NMR (62.5 MHz, DMSO-CD<sub>3</sub>OD (1/2)) δ (ppm):

173 (3  $\text{CH}_3\text{CO}$ ), 170 ( $\text{NCOAr}$ ), 158 ( $\text{ArC-3}$ ), 137 ( $\text{ArC-1}$ ), 131 ( $\text{ArC-5}$ ), 121 ( $\text{ArC-6}$ ), 119 ( $\text{ArC-4}$ ), 115 ( $\text{ArC-2}$ ), 103 ( $\text{C-1}\beta^{\text{II,III,IV}}$ ), 96 ( $\text{C-1}\beta^{\text{I}}$ ), 92 ( $\text{C-1}\alpha^{\text{I}}$ ), 82 – 50 (20 C of the sugars,  $\text{C}\equiv\text{C}$  and  $\text{CH}_2\text{-OAr}$ ), 33-16 (7  $\text{CH}_2$  and 3  $\text{CH}_3\text{CO}$ ), 15 ( $\text{CH}_3$ )

5 Mass spectrum:

Negative ESI  $m/z = 1137.1$   $[\text{M}-\text{Na}]^-$

10 **2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(*N*-2-(undec-4Z-  
enyloxy)benzoyl)amino- $\beta$ -D-glucopyranosyl)- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-6-O-sulfo-D-glucopyranose, sodium salt (**8**)**



15 10 mg of **CO-IV(NH<sub>2</sub>S)** (11  $\mu\text{mol}$ ) are dissolved in 100  $\mu\text{L}$  of water and 250  $\mu\text{L}$  of DMF. 2 mg of sodium hydrogen carbonate (24  $\mu\text{mol}$ ) and then 15  $\mu\text{L}$  of a solution of **31** in THF at a concentration of 115 mg/mL (6  $\mu\text{mol}$ ) are then added. The reaction medium is heated to 60°C and 105  $\mu\text{L}$  of the solution of chloride and 6 mg of sodium hydrogen carbonate are added in seven portions over 18 hours. After concentrating, the residue is purified by placing it in DCM/methanol (5/1) on a column of silica, while diluting it greatly, in  
20 order to remove the lipid chain. The elution is then performed with E/M/W (9/2/1). 6.2 mg of a white solid are thus isolated, i.e. a yield of 48% (but a conversion of only 50%).

$^1\text{H}$  NMR (400 MHz, DMSO- $\text{CD}_3\text{OD}$  (1/2))  $\delta$  (ppm):

7.99 (dd, 1 H,  $\text{ArH-6}$ ,  $J_{6,5}$  7.5 Hz and  $J_{6,4}$  1.8 Hz), 7.55 (ddd, 1 H,  $\text{ArH-4}$ ,  $J_{4,3}$  8.3 Hz and  $J_{4,5}$  7.8 Hz), 7.20  
25 (d, 1 H,  $\text{ArH-3}$ ), 7.10 (dd, 1H,  $\text{ArH-5}$ ), 5.52 (m, 2 H,  $\text{CH=CH}$ ), 5.06 (d, 0.7 H,  $\text{H-1}\alpha^{\text{I}}$ ,  $J_{1,2}$  3.0 Hz), 4.70-  
4.60-4.53 (4 d superimposed, 3.6 H,  $\text{H-1}\beta^{\text{I,II,III,IV}}$ ), 4.20-3.40 (m, 25 H, other Hs of the sugars and  $\text{CH}_2\text{-OAr}$ ), 2.33-2.11 (2 m, 4 H,  $\text{CH}_2\text{-CH=CH-CH}_2$ ), 2.03-2.01-2.00 (3 s, 9 H, 3  $\text{CH}_3\text{CO}$ ), 2.05 (m, 2 H,  $\text{ArO-CH}_2\text{-CH}_2\text{-CH}_2$ ), 1.50-1.20 (m, 8 H, 4  $\text{CH}_2$ ), 0.94 (t, 3 H,  $\text{CH}_3$ ,  $J$  6.8 Hz)


$^{13}\text{C}$  NMR (62.5 MHz, DMSO- $\text{CD}_3\text{OD}$  (1/2))  $\delta$  (ppm):

172 (3  $\text{CH}_3\text{CO}$ ), 171 ( $\text{NCOAr}$ ), 158 ( $\text{ArC-1}$ ), 133 ( $\text{ArC-4}$ ,  $\text{CH=CH}$ ), 129 ( $\text{ArC-6}$ ), 122 ( $\text{ArC-5}$ ), 114 ( $\text{ArC-3}$ ), 103 ( $\text{C-1}\beta^{\text{II,III,IV}}$ ), 96 ( $\text{C-1}\beta^{\text{I}}$ ), 92 ( $\text{C-1}\alpha^{\text{I}}$ ), 82-50 (all the other Cs of the sugars and  $\text{CH}_2\text{OAr}$ ), 33-24 (7  $\text{CH}_2$  and 3  $\text{CH}_3\text{CO}$ ), 15 ( $\text{CH}_3$ )

Mass spectrum:

Negative ESI  $m/z = 1139.5$  [ $\text{M}-\text{Na}$ ]-

**2-acetamido-4-O-{2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(*N*-4-(undec-4Z-enoxy)benzoyl)amino- $\beta$ -D-glucopyranosyl)- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-6-O-sulfo-D-glucopyranose, sodium salt (**9**)**



10 mg of **CO-IV(NH<sub>2</sub>,S)** (11  $\mu\text{mol}$ ) are dissolved in 100  $\mu\text{L}$  of water and 250  $\mu\text{L}$  of DMF. 2 mg of sodium hydrogen carbonate (24  $\mu\text{mol}$ ) and then 15  $\mu\text{L}$  of a solution of **35** in THF at a concentration of 115 mg/mL (6  $\mu\text{mol}$ ) are then added. The reaction medium is heated to 60°C and 105  $\mu\text{L}$  of the solution of chloride and 6 mg of sodium hydrogen carbonate are added in seven portions over 17 hours. After concentrating, the residue is purified by placing it in DCM/methanol (5/1) on a column of silica, while diluting it greatly, in order to remove the lipid chain. The elution is then performed with E/MW (9/2/1). 5.2 mg of a white solid are thus isolated, i.e. a yield of 40% (but a conversion of only 60%).

$^1\text{H}$  NMR (400 MHz, DMSO- $\text{CD}_3\text{OD}$  (1/2))  $\delta$  (ppm):

7.89 (d, 2 H,  $\text{ArH-2}$  and  $\text{ArH-6}$ ,  $J_{2,3} \approx J_{6,5}$  8.8 Hz), 7.04 (d, 2 H,  $\text{ArH-3}$  and  $\text{ArH-5}$ ), 5.48 (m, 2 H,  $\text{CH=CH}$ ), 5.05 (d, 0.6 H,  $\text{H-1}\alpha^{\text{I}}$ ,  $J_{1\alpha,2}$  3.1 Hz), 4.69-4.55-4.50 (4 d superimposed, 3.6 H,  $\text{H-1}\beta^{\text{I,II,III,IV}}$ ), 4.30-3.40 (m, 23 H, other Hs of the sugars), 4.10 (t,  $\text{CH}_2\text{-OAr}$ ,  $J$  6.3 Hz), 2.28-2.09 (2 m, 4 H,  $\text{CH}_2\text{-CH=CH-CH}_2$ ), 2.02-1.99-1.97 (3 s, 9 H, 3  $\text{CH}_3\text{CO}$ ), 1.89 (m, 2 H,  $\text{ArO-CH}_2\text{-CH}_2\text{-CH}_2$ ), 1.45-1.25 (m, 8 H, 4  $\text{CH}_2$ ), 0.93 (t, 3 H,  $\text{CH}_3$ ,  $J$  7.0 Hz)


$^{13}\text{C}$  NMR (62.5 MHz, DMSO- $\text{CD}_3\text{OD}$  (1/2))  $\delta$  (ppm):

172 (3  $\text{CH}_3\text{CO}$ ), 169 ( $\text{NCOAr}$ ), 163 ( $\text{ArC-1}$ ), 132-130-129 ( $\text{ArC-2}$ ,  $\text{ArC-6}$ ,  $\text{CH=CH}$ ), 115 ( $\text{ArC-3}$ ,  $\text{ArC-5}$ ), 103 ( $\text{C-1}\beta^{\text{II,III,IV}}$ ), 97 ( $\text{C-1}\beta^{\text{I}}$ ), 92 ( $\text{C-1}\alpha^{\text{I}}$ ), 83-50 (all the other Cs of the sugars and  $\text{CH}_2\text{OAr}$ ), 33-23 (7  $\text{CH}_2$  and 3  $\text{CH}_3\text{CO}$ ), 15 ( $\text{CH}_3$ )

Mass spectrum:

Negative ESI  $m/z = 1139.5$   $[\text{M}-\text{Na}]^-$

**2-acetamido-4-O-[2-acetamido-4-O-[2-acetamido-4-O-[2-acetamido-2-deoxy-4-O-(2-deoxy-2-(N-3-(undec-4Z-enyloxy)benzoyl)amino- $\beta$ -D-glucopyranosyl)- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy- $\beta$ -D-glucopyranosyl]-2-deoxy-6-O-( $\alpha$ -L-fucopyranosyl)-D-glucopyranose**  
**(10)**




The fucosyl pentamer **CO-V(NH<sub>2</sub>, Fuc)** (7.3 mg, 6.4  $\mu\text{mol}$ ) is dissolved in  $\text{H}_2\text{O}$  (140  $\mu\text{L}$ ), followed by addition of DMF (350  $\mu\text{L}$ ) and the mixture is brought to 30°C. Dowex 1x2-100 resin ( $\text{HCO}_3^-$ ) is then added, followed by addition of a solution (THF, 110  $\mu\text{L}$ ) of the acid chloride **19** (6 mg). The reaction mixture is stirred for 24 hours, during which time three further additions of resin and of acid chloride solution are made. The reaction medium is then diluted in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (1/1, 2 mL), heated to 56°C, and the supernatant is then filtered through cotton wool. The resin beads and the walls of the flask are extracted several times at 56°C with  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (4/1, 7/3, 3/2, 1/1, 2/3, 3/7 and 1/4, 2 mL each). The various fractions are passed through a Dowex 50x8-100 resin ( $\text{H}^+$ ), and then pooled and concentrated. The residue is successively washed with EtOAc (3 x 1 mL) and then  $\text{H}_2\text{O}$  (3 x 1 mL), and then redissolved in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (1/1, 10 mL) by heating to 56°C, and then by sonication. The solution is then freeze-dried, and the expected product is obtained in the form of a white powder (2.5 mg, 28%).

The starting material retained on the acid resin is then eluted (2.3 mg, 31%) using aqueous ammonia solution (H<sub>2</sub>O, 2%).

<sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>/D<sub>2</sub>O 20/1) δ (ppm):

5 7.43 – 7.30 (m, 3 H, ArH-2, ArH-6 and ArH-5); 7.05 (m, 1 H, ArH-4); 5.45 – 5.32 (m, 2 H, CH=CH); 4.84 (d, 0.8H, *J*<sub>1,2</sub> = 1.9 Hz, H-1α<sup>I</sup>); 4.66 (d, 0.8H, *J*<sub>1,2</sub> < 1.0 Hz, H-1Fuc-GlcNAcα), 4.65 (d, 0.2H, *J*<sub>1,2</sub> < 1.0 Hz, H-1Fuc-GlcNAcβ), 4.52 (d, H, *J* = 8.5 Hz, H-1β<sup>IV</sup>), 4.45 / 4.35 / 4.33 (4d, 4H, *J* = 8.5 Hz, H-1β<sup>II-IV</sup>), 4.42 (d, 0.2H, *J* = 7.0 Hz, H-1β<sup>I</sup>); 3.99 (t, 2H, *J* = 6.1 Hz, ArOCH<sub>2</sub>-CH<sub>2</sub>), 3.88 (dt, 1H, H-5Fuc), 3.78 – 3.05 (m, 33H, other sugar Hs), 2.17 (dt, 2H, *J* = 6.0 and *J* = 6.8 Hz, CH<sub>2</sub>-CH=CH), 1.99 (dt, 2H, *J* = 5.9 and *J* = 6.2 Hz, CH=CH-CH<sub>2</sub>), 1.82 / 1.81 / 1.81 / 1.79 (4s, 12H, 4 COCH<sub>3</sub>), 1.80 – 1.72 (m, 2 H, ArOCH<sub>2</sub>-CH<sub>2</sub>), 1.31 – 1.15 (m, 8 H, 4 CH<sub>2</sub>), 1.08 (d, 0.6H, *J*<sub>5,6</sub> = 6.9 Hz, H-6Fuc-GlcNAcβ), 1.05 (d, 2.4H, *J*<sub>5,6</sub> = 6.5 Hz, H-6Fuc-GlcNAcα), 0.82 (t, 3 H, CH<sub>3</sub>, *J* = 6.5 Hz).

15 **methyl 3-(undec-4Z-enyloxy)benzoate (15)**



850 mg of 14 (6.15 mmol) and 900 mg of K<sub>2</sub>CO<sub>3</sub> (6.51 mmol) are added to 1.7 g of 13 (6.07 mmol) in anhydrous DMF (20 mL). After reaction for 4 hours at 90°C, the reaction medium is concentrated, taken up in DCM and then washed with water. 1.87 g of a yellow oil are obtained, and are chromatographed on silica gel in pentane/ethyl acetate (50/1). 1.37 g of a yellow oil are isolated, i.e. a yield of 76%.

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ (ppm):

25 7.60 (ddd, 1 H, ArH-6, *J*<sub>6,5</sub> 8.0 Hz and *J*<sub>6,4</sub> ≈ *J*<sub>6,2</sub> 0.5 Hz), 7.52 (dd, 1 H, ArH-2, *J*<sub>2,4</sub> 3.0 Hz), 7.31 (dd, 1 H, ArH-5, *J*<sub>5,4</sub> 8.0 Hz), 7.07 (ddd, 1 H, ArH-4), 5.38 (m, 2 H, CH=CH), 3.98 (t, 2 H, CH<sub>2</sub>-OAr, *J* 6.3 Hz), 3.89 (s, 3 H, OCH<sub>3</sub>), 2.22-1.99 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.83 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH=CH, *J* 6.8 Hz), 1.55-1.20 (m, 8 H, 4 CH<sub>2</sub>), 0.84 (t, 3 H, CH<sub>3</sub>, *J* 7.5 Hz)

30 <sup>13</sup>C NMR (62.5 MHz, CDCl<sub>3</sub>) δ (ppm):

131-129-128 (C-5, CH=CH), 122 (C-6), 120 (C-4), 115 (C-2), 66 (CH<sub>2</sub>-OAr), 52 (CH<sub>3</sub>O), 32-22 (7 CH<sub>2</sub>), 14 (CH<sub>3</sub>)

Mass spectrum:

Positive ESI  $m/z = 327.2$   $[M + Na]^+$

High res. Calc. for  $C_{19}H_{28}O_3Na$ : 327.193614, Found : 327.193200

5

Elemental analysis:


	Calc.	Found
C	74.96	74.68
H	9.27	9.37
O	15.77	15.79

10

Infrared ( $cm^{-1}$ ): 2970-2950-2927-2858-1726-1586-1446-1288-1228-756

### 3-(undec-4Z-enoxy)benzyl alcohol (**16**)

15



35 mg of lithium aluminum hydride (922  $\mu$ mol) are added, at 0°C, to 140 mg of **15** (460  $\mu$ mol) in ether (3 mL). After reaction for 1 hour 30 minutes, the reaction medium is diluted with ether and hydrolyzed with two drops of water. After filtering through Celite, drying over  $Na_2SO_4$  and concentrating, 127 mg of a colorless oil are isolated, i.e. a yield of 99%.

20

$^1H$  NMR (250 MHz,  $CDCl_3$ )  $\delta$  (ppm):

7.18 (dd, 1 H, ArH-5,  $J_{5,6}$  8.0 Hz and  $J_{5,4}$  8.3 Hz), 6.84 (m, 2 H, ArH-2 and ArH-4), 6.75 (dd, 1 H, ArH-4,  $J_{4,2}$  2.9 Hz), 5.32 (m, 2 H, CH=CH), 4.58 (s, 2 H, CH<sub>2</sub>OH), 3.89 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 2.16-1.95 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.76 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH=CH,  $J$  6.8 Hz), 1.45-1.18 (m, 8 H, 4 CH<sub>2</sub>), 0.84 (t, 3 H, CH<sub>3</sub>,  $J$  6.3 Hz)

25

$^{13}C$  NMR (62.5 MHz,  $CDCl_3$ )  $\delta$  (ppm):

159 (C-3), 142 (C-1), 131-130-128 (C-5, CH=CH), 119 (C-6), 114 (C-4), 113 (C<sub>2</sub>), 67 (CH<sub>2</sub>-OAr), 65 (CH<sub>2</sub>OH), 32-23 (7 CH<sub>2</sub>), 14 (CH<sub>3</sub>)

30


Mass spectrum:

Positive ESI  $m/z = 299.2$   $[M + Na]^+$

High res. Calc. for  $C_{18}H_{28}O_2Na$ : 299.198700, Found : 299.199250

Infrared ( $cm^{-1}$ ): 3329, 3005, 2940, 2925, 2855, 1669, 1602, 1452, 1264

5 **3-(undec-4Z-enyloxy)benzaldehyde (17)**



10 mL of anhydrous DCM and then 190 mg of PCC (881  $\mu$ mol) are added under argon to 120 mg of alcohol **16** (434  $\mu$ mol) dried by coevaporation with toluene. The reaction is heated to the reflux point of the DCM for 1 hour. After cooling, the reaction medium is diluted with ether and filtered through Florisil. After concentrating, 118 mg of a yellow oil are obtained, i.e. a yield of 99%.

$^1H$  NMR (250 MHz,  $CDCl_3$ )  $\delta$  (ppm):

15 9.95 (s, 1 H,  $\underline{CHO}$ ), 7.42 (m, 2 H,  $\underline{ArH-6}$  and  $\underline{ArH-5}$ ), 7.36 (d, 1 H,  $\underline{ArH-2}$ ,  $J_{2,4}$  2.9 Hz), 7.15 (m, 1 H,  $\underline{ArH-4}$ ), 5.39 (m, 2 H,  $\underline{CH=CH}$ ), 3.99 (t, 2 H,  $\underline{CH_2-OAr}$ ,  $J$  6.3 Hz), 2.21-1.99 (2 m, 4 H,  $\underline{CH_2-CH=CH-CH_2}$ ), 1.84 (tt, 2 H,  $\underline{ArO-CH_2-CH_2-CH_2-CH=CH}$ ,  $J$  6.8 Hz), 1.40 – 1.15 (m, 8 H, 4  $\underline{CH_2}$ ), 0.84 (t, 3 H,  $\underline{CH_3}$ ,  $J$  6.6 Hz)

$^{13}C$  NMR (62.5 MHz,  $CDCl_3$ )  $\delta$  (ppm):

20 192 ( $\underline{CHO}$ ), 160 ( $\underline{C-3}$ ), 138 ( $\underline{C-1}$ ), 131-130-128 ( $\underline{C-5}$ ,  $\underline{CH=CH}$ ), 123 ( $\underline{C-6}$ ), 122 ( $\underline{C-4}$ ), 113 ( $\underline{C-2}$ ), 67 ( $\underline{CH_2-OAr}$ ), 52 ( $\underline{CH_3O}$ ), 32-23 (7  $\underline{CH_2}$ ), 14 ( $\underline{CH_3}$ )

Mass spectrum:

25 Chemical ionization (CI) 1% solution in DCM


A fine desorption peak

$M + 1 = 275$

Infrared ( $cm^{-1}$ ): 3005-2940-2927-2855-2723-1700-1599-1452-1263-787

30

**3-(undec-4Z-enyloxy)benzoic acid (18)**



4 mL of 1 N sodium hydroxide solution (4.0 mmol) are added portionwise to 1.14 g of **15** (3.74 mmol) in methanol (30 mL). The solution is refluxed overnight. A further 4 mL of 1 N sodium hydroxide solution are added, and the mixture is refluxed for a further 1 hour 30 minutes. After evaporating off the solvent, the reaction medium is acidified with 0.5 N HCl and extracted with DCM. 1.04 g of a yellow oil are obtained, i.e. a yield of 96%.

$^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

10.00-9.00 (bd, 1 H,  $\text{CO}_2\text{H}$ ), 7.69 (d, 1 H, ArH-6,  $J_{6,5}$  7.8 Hz), 7.60 (d, 1 H, ArH-2,  $J_{2,4}$  2.4 Hz), 7.35 (dd, 1 H, ArH-5,  $J_{5,4}$  8.3 Hz), 7.14 (dd, 1 H, ArH-4), 5.40 (m, 2 H,  $\text{CH}=\text{CH}$ ), 4.00 (t, 2 H,  $\text{CH}_2\text{-OAr}$ ,  $J$  6.3 Hz), 2.21-2.00 (2 m, 4 H,  $\text{CH}_2\text{-CH}=\text{CH-CH}_2$ ), 1.85 (tt, 2 H,  $\text{ArO-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}=\text{CH}$ ,  $J$  6.8 Hz), 1.35-1.05 (m, 8 H, 4  $\text{CH}_2$ ), 0.85 (t, 3 H,  $\text{CH}_3$ ,  $J$  6.5 Hz)

$^{13}\text{C}$  NMR (50 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

172 ( $\text{C}=\text{O}$ ), 159 ( $\text{C}-3$ ), 131-130-130-128 ( $\text{C}-1$ ,  $\text{C}-5$ ,  $\text{CH}=\text{CH}$ ), 121-122 ( $\text{C}-4$ ,  $\text{C}-6$ ), 115 ( $\text{C}-2$ ), 67 ( $\text{CH}_2\text{-OAr}$ ), 32-23 (7  $\text{CH}_2$ ), 14 ( $\text{CH}_3$ )

Mass spectrum:

Negative ESI  $m/z$  = 289.1  $[\text{M-H}]^-$

High res. Calc. for  $\text{C}_{18}\text{H}_{25}\text{O}_3$  : 289.180370, Found : 289.180730


Elemental analysis:

	Calc.	Found
C	74.45	74.29
H	9.02	9.01

Infrared ( $\text{cm}^{-1}$ ): 2970-2950-2925-2854-1695-1585-1286-757

**3-(undec-4Z-enyloxy)benzoic acid (**15**)**






1 mL of oxalyl chloride (11.5 mmol) and two drops of anhydrous DMF are added under argon to 100 mg of toluene-dried **18** (345  $\mu$ mol) dissolved in 20 mL of anhydrous DCM. The medium is stirred at room temperature for two hours, and then concentrated to give 106 mg of the expected chloride in the form of a yellow oil, i.e. a yield of 99%.

$^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

7.71 (ddd, 1 H, ArH-6,  $J_{6,5}$  8.3 Hz,  $J_{6,4}$  2.4 Hz and  $J_{6,2}$  0.9 Hz), 7.57 (dd, 1 H, ArH-2,  $J_{2,4}$  1.6 Hz), 7.39 (dd, 1 H, ArH-5,  $J_{5,4}$  8.3 Hz), 7.20 (ddd, 1 H, ArH-4), 5.40 (m, 2 H, CH=CH), 3.99 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 2.23-2.00 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.85 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH=CH,  $J$  7.0 Hz), 1.28-1.15 (m, 8 H, 4 CH<sub>2</sub>), 0.85 (t, 3 H, CH<sub>3</sub>,  $J$  6.5 Hz)

#### methyl 3-(undecyloxy)benzoate (**21**)



350 mg of **14** (2.30 mmol) and 330 mg of  $\text{K}_2\text{CO}_3$  (2.39 mmol) are added to 554 mg of 1-bromoundecane (2.35 mmol) in anhydrous DMF (7 mL). After reaction for 16 hours at 90°C, the reaction medium is concentrated, taken up in DCM and then washed with water. 607 mg of a yellow oil are obtained, and are chromatographed on silica gel in pentane/ethyl acetate (60/1). 579 mg of expected coupling product are isolated in the form of a yellow oil, i.e. a yield of 82%.

$^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

7.62 (m, 1 H, ArH-6), 7.55 (m, 1 H, ArH-2), 7.34 (dd, 1 H, ArH-5,  $J_{5,4}$  8.1 Hz and  $J_{5,6}$  7.7 Hz), 7.10 (ddd, 1 H, ArH-4,  $J_{4,6}$  2.8 Hz and  $J_{4,2}$  0.8 Hz), 4.00 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.6 Hz), 3.92 (s, 3 H, OCH<sub>3</sub>), 1.80 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>,  $J$  6.6 Hz and  $J$  6.4 Hz), 1.52-1.20 (m, 16 H, 8 CH<sub>2</sub>), 0.89 (t, 3 H, CH<sub>3</sub>,  $J$  6.7 Hz)

$^{13}\text{C}$  NMR (62.5 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

167 (CO<sub>2</sub>CH<sub>3</sub>), 159 (C-3), 131 (C-1), 129 (C-5), 122 (C-6), 120 (C-4), 115 (C-2), 68 (CH<sub>2</sub>-OAr), 52 (CH<sub>3</sub>O), 32-23 (9 CH<sub>2</sub>), 14 (CH<sub>3</sub>)


Mass spectrum:

5 Positive ESI  $m/z = 329.2$  [M + Na]<sup>+</sup>

High res. Calc. for C<sub>19</sub>H<sub>30</sub>O<sub>3</sub>Na: 329.209264, Found : 329.207940

Infrared (cm<sup>-1</sup>): 2950-2925-2854-1727-1586-1446-1287-1228-756

10 **3-(undecyloxy)benzoic acid (22)**



600  $\mu$ L of 1 N sodium hydroxide solution (600  $\mu$ mol) are added portionwise to 112 mg of 21 (366  $\mu$ mol) in  
15 methanol (4 mL). The solution is refluxed for two hours. After evaporating off the solvent, the reaction  
medium is acidified with 0.5 N HCl and extracted with DCM. 107 mg of the expected acid are obtained in  
the form of a white solid, i.e. a yield of 99%.

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm):

20 7.70 (d, 1 H, ArH-6,  $J_{6,5}$  7.8 Hz), 7.62 (m, 1 H, ArH-2), 7.38 (dd, 1 H, ArH-5,  $J_{5,4}$  8.0 Hz), 7.16 (dd, 1 H,  
ArH-4,  $J_{4,2}$  2.1 Hz), 4.02 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.5 Hz), 1.99 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>,  $J$  6.6 Hz), 1.55-1.20  
(m, 16 H, 8 CH<sub>2</sub>), 0.89 (t, 3 H, CH<sub>3</sub>,  $J$  6.5 Hz)

<sup>13</sup>C NMR (62.5 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm):

25 171 (CO<sub>2</sub>H), 159 (C-3), 130 (C-1), 129 (C-5), 122 (C-6), 121 (C-4), 115 (C-2), 68 (CH<sub>2</sub>-OAr), 32-23 (9  
CH<sub>2</sub>), 14 (CH<sub>3</sub>)

Mass spectrum:


Negative ESI  $m/z = 291.2$  [M-H]<sup>-</sup>

30 High res. Calc. for C<sub>18</sub>H<sub>27</sub>O<sub>3</sub> : 291.196020, Found : 291.196560

Infrared (cm<sup>-1</sup>): 2950-2920-2850-2700-2400-1680-1603-1455-1420-1312-1247-757


Melting point: 88°C

**3-(undecanyloxy)benzoyl chloride (23)**



1 mL of oxalyl chloride (11.5 mmol) and two drops of anhydrous DMF are added under argon to 93 mg of toluene-dried acid 22 (318  $\mu$ mol) dissolved in 20 mL of anhydrous DCM. The medium is stirred at room temperature for two hours, and then concentrated to give 99 mg of a yellow oil, i.e. a yield of 99%.

**methyl 3-(undec-4-ynyloxy)benzoate (25)**



325 mg of 14 (2.14 mmol) and 300 mg of  $K_2CO_3$  (2.17 mmol) are added to 600 mg of 24 (2.16 mmol) in anhydrous DMF (7 mL). After reaction for 6 hours at 90°C, the reaction medium is concentrated, washed with DCM and then taken up in water. 639 mg of a yellow oil are obtained, and are chromatographed on silica gel in pentane/ethyl acetate (50/1). 429 mg of a yellow oil are isolated, i.e. a yield of 66%.

$^1H$  NMR (200 MHz,  $CDCl_3$ )  $\delta$  (ppm):

7.63 (m, 1 H, ArH-6), 7.57 (m, 1 H, ArH-2), 7.31 (dd, 1 H, ArH-5,  $J_{5,4}$  8.1 Hz and  $J_{5,6}$  7.8 Hz), 7.11 (ddd, 1 H, ArH-4,  $J_{4,6}$  2.4 Hz and  $J_{4,2}$  0.8 Hz), 4.11 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.2 Hz), 3.92 (s, 3 H, OCH<sub>3</sub>), 2.39-2.15 (2 m, 4 H, CH<sub>2</sub>-C≡C-CH<sub>2</sub>), 1.98 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-C≡C,  $J$  6.5 Hz), 1.52-1.23 (m, 8 H, 4 CH<sub>2</sub>), 0.88 (t, 3 H, CH<sub>3</sub>,  $J$  6.7 Hz)

$^{13}C$  NMR (62.5 MHz,  $CDCl_3$ )  $\delta$  (ppm):

167 (C=O<sub>2</sub>CH<sub>3</sub>), 159 (C-3), 131 (C-1), 129 (C-5), 122 (C-6), 120 (C-4), 115 (C-2), 81-79 (C≡C), 67 (CH<sub>2</sub>-OAr), 52 (CH<sub>3</sub>O), 31-15 (7 CH<sub>2</sub>), 14 (CH<sub>3</sub>)


Mass spectrum:

Positive ESI  $m/z$  = 325.1 [M + Na]<sup>+</sup>

High res. Calc. for  $C_{19}H_{26}O_3Na$ : 325.177964, Found : 325.178070

Infrared ( $cm^{-1}$ ): 2950-2931-2857-1726-1586-1446-1288-1228-756

5 **3-(undec-4-ynyloxy)benzoic acid (26)**



300  $\mu$ L of 1 N sodium hydroxide solution (300  $\mu$ mol) are added portionwise to 48 mg of **25** (157  $\mu$ mol) in  
10 methanol (2 mL). The solution is refluxed for 1 hour 30 minutes. After evaporating off the solvent, the  
reaction medium is acidified with 0.5 N HCl and extracted with DCM. 45 mg of a pale yellow oil are  
obtained, i.e. a yield of 99%.

$^1H$  NMR (250 MHz,  $CDCl_3$ )  $\delta$  (ppm):

15 11.00-10.00 (bd, 1 H,  $CO_2H$ ), 7.72 (d, 1 H,  $ArH$ -6,  $J_{6,5}$  7.7 Hz), 7.64 (m, 1 H,  $ArH$ -2), 7.38 (dd, 1 H,  $ArH$ -5,  
 $J_{5,4}$  8.1 Hz), 7.17 (dd, 1 H,  $ArH$ -4,  $J_{4,2}$  2.7 Hz), 4.13 (t, 2 H,  $CH_2$ -OAr,  $J$  6.1 Hz), 2.39-2.15 (2 m, 4 H,  
 $CH_2$ -C $\equiv$ C- $CH_2$ ), 1.99 (tt, 2 H, ArO- $CH_2$ - $CH_2$ -C $\equiv$ C,  $J$  6.5 Hz), 1.50-1.20 (m, 8 H, 4  $CH_2$ ), 0.88 (t, 3 H,  
 $CH_3$ ,  $J$  6.7 Hz)

20  $^{13}C$  NMR (62.5 MHz,  $CDCl_3$ )  $\delta$  (ppm):

172 ( $CO_2H$ ), 159 ( $C$ -3), 131 ( $C$ -1), 129 ( $C$ -5), 123 ( $C$ -6), 121 ( $C$ -4), 115 ( $C$ -2), 81-79 ( $C\equiv C$ ), 67 ( $CH_2$ -OAr),  
31-15 (7  $CH_2$ ), 14 ( $CH_3$ )


Mass spectrum:

25 Negative ESI  $m/z$  = 287.1 [M-H] $^-$

High res. Calc. for  $C_{18}H_{23}O_3$  : 287.164719, Found : 287.164820


Infrared ( $cm^{-1}$ ): 2954-2929-2855-2700-2400-1690-1592-1452-1414-1288-1247-756

30 **3-(undec-4Z-ynyloxy)benzoyl chloride (27)**



850  $\mu$ L of oxalyl chloride (9.74 mmol) and two drops of anhydrous DMF are added under argon to 80 mg of toluene-dried acid **26** (278  $\mu$ mol) dissolved in 17 mL of anhydrous DCM. The medium is stirred at room temperature for two hours, and then concentrated to give 85 mg of a yellow oil, i.e. a yield of 99%.

#### methyl 2-(undec-4Z-enyloxy)benzoate (**29**)



88 mg of **28** (578  $\mu$ mol) and 77 mg of  $K_2CO_3$  (557  $\mu$ mol) are added to 140 mg of **13** (500  $\mu$ mol) in anhydrous DMF (2 mL). After reaction for 8 hours at 90°C, the reaction medium is concentrated, taken up in DCM and then washed with water. 137 mg of a yellow oil are obtained, and are chromatographed on silica gel in pentane/ethyl acetate (40/1). 100 mg of a yellow oil are isolated, i.e. a yield of 66%.

$^1H$  NMR (250 MHz,  $CDCl_3$ )  $\delta$  (ppm):

7.79 (dd, 1 H, ArH-6,  $J_{6,5}$  8.1 Hz and  $J_{6,4}$  1.9 Hz), 7.43 (ddd, 1 H, ArH-4,  $J_{4,3}$  8.5 Hz,  $J_{4,5}$  7.3 Hz), 6.94 (m, 2 H, ArH-5 and ArH-3), 5.40 (m, 2 H, CH=CH), 4.02 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 3.89 (s, 3 H, OCH<sub>3</sub>), 2.28-2.01 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>,  $J$  6.6 Hz), 1.89 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>,  $J$  6.6 Hz), 1.50-1.16 (m, 8 H, 4 CH<sub>2</sub>), 0.86 (t, 3 H, CH<sub>3</sub>,  $J$  6.6 Hz)

$^{13}C$  NMR (62.5 MHz,  $CDCl_3$ )  $\delta$  (ppm):

167 (C=O), 158 (C-2), 133 (C-4), 131 (CH=CH), 128 (C-6), 120 (C-1), 119 (C-5), 113 (C-3), 68 (CH<sub>2</sub>-OAr), 52 (CH<sub>3</sub>O), 32-22 (7 CH<sub>2</sub>), 14 (CH<sub>3</sub>)


Mass spectrum:

Positive ESI  $m/z$  = 327.2 [M + Na]<sup>+</sup>

High res. Calc. for C<sub>19</sub>H<sub>28</sub>O<sub>3</sub>Na: 327.193914, Found : 327.192560

Infrared (cm<sup>-1</sup>): 3000-2962-2925-2855-1734-1601-1491-1456-1305-1250-754

**2-(undec-4Z-enyloxy)benzoic acid (30)**



5

500  $\mu$ L of 1 N sodium hydroxide solution (500  $\mu$ mol) are added portionwise to 80 mg of **29** (263  $\mu$ mol) in methanol (3 mL). The solution is refluxed for 24 hours. After evaporating off the solvent, the reaction medium is acidified with 0.5 N HCl and extracted with DCM. 76 mg of a yellow oil are obtained, i.e. a yield of 99%.

10

$^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

12.00-10.00 (bd, 1 H,  $\text{CO}_2\text{H}$ ), 8.16 (dd, 1 H, ArH-6,  $J_{6,5}$  7.8 Hz and  $J_{6,4}$  1.9 Hz), 7.54 (ddd, 1 H, ArH-4,  $J_{4,3}$  8.4 Hz and  $J_{4,5}$  7.6 Hz), 7.10 (ddd, 1 H, ArH-5,  $J_{5,3}$  0.8 Hz), 7.03 (dd, 1 H, ArH-3), 5.40 (m, 2 H,  $\text{CH}=\text{CH}$ ), 4.24 (t, 2 H,  $\text{CH}_2\text{-OAr}$ ,  $J$  6.4 Hz), 2.25 (m, 2 H,  $\text{CH}_2\text{-CH}=\text{CH-CH}_2$ ), 1.97 (m, 4 H,  $\text{ArO-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}=\text{CH-CH}_2$ ), 1.35-1.10 (m, 8 H, 4  $\text{CH}_2$ ), 0.84 (t, 3 H,  $\text{CH}_3$ ,  $J$  6.6 Hz)

15

$^{13}\text{C}$  NMR (62.5 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

165 ( $\text{C}=\text{O}$ ), 157 ( $\text{C}-2$ ), 135 ( $\text{C}-4$ ), 134-132 ( $\text{CH}=\text{CH}$ ), 127 ( $\text{C}-6$ ), 122 ( $\text{C}-5$ ), 117 ( $\text{C}-1$ ), 112 ( $\text{C}-3$ ), 69 ( $\text{CH}_2\text{-OAr}$ ), 32-22 (7  $\text{CH}_2$ ), 14 ( $\text{CH}_3$ )

20


Mass spectrum:

Negative ESI  $m/z$  = 289.2  $[\text{M-H}]^-$

High res. Calc. for  $\text{C}_{18}\text{H}_{25}\text{O}_3$ : 289.180370, Found : 289.179060

25

**2-(undec-4Z-enyloxy)benzoyl chloride (31)**




800  $\mu$ L of oxalyl chloride (9.17 mmol) and two drops of anhydrous DMF are added under argon to 76 mg of toluene-dried acid **30** (262  $\mu$ mol) dissolved in 15 mL of anhydrous DCM. The medium is stirred at room temperature for two hours and then concentrated to give 80 mg of a yellow oil, i.e. a yield of 99%.

5  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

7.97 (dd, 1 H, ArH-6,  $J_{6,5}$  7.9 Hz and  $J_{6,4}$  1.7 Hz), 7.46 (m, 1 H, ArH-4), 6.90 (m, 2H, ArH-5 and ArH-3), 5.30 (m, 2 H, CH=CH), 3.95 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 2.20-1.90 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.79 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH=CH,  $J$  6.6 Hz), 1.20-1.09 (m, 8 H, 4 CH<sub>2</sub>), 0.76 (t, 3 H, CH<sub>3</sub>,  $J$  6.7 Hz)

10 methyl 4-(undec-4Z-enyloxy)benzoate (**33**)



90 mg of **32** (590  $\mu$ mol) and 81 mg of  $\text{K}_2\text{CO}_3$  (590  $\mu$ mol) are added to 150 mg of **13** (535  $\mu$ mol) in  
15 anhydrous DMF (2 mL). After reaction for 7 hours at 90°C, the reaction medium is concentrated, taken up in DCM and then washed with water. 163 mg of a yellow oil are obtained, and are chromatographed on silica gel in pentane/ethyl acetate (80/1). 129 mg of the expected coupling product are isolated in the form of a yellow oil, i.e. a yield of 79%.

20  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

7.97 (d, 2 H, ArH-2 and ArH-6,  $J_{6,5} \approx J_{2,3}$  8.8 Hz), 6.89 (d, 2 H, ArH-3 and ArH-5), 5.39 (m, 2 H, CH=CH), 3.99 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 3.88 (s, 3 H, OCH<sub>3</sub>), 2.22-2.00 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.84 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>,  $J$  6.8 Hz), 1.40-1.12 (m, 8 H, 4 CH<sub>2</sub>), 0.85 (t, 3 H, CH<sub>3</sub>,  $J$  6.6 Hz)

25  $^{13}\text{C}$  NMR (62.5 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm): 167 (C=O), 163 (C-4), 131 (C-2 and C-6), 130-128 (CH=CH), 122 (C-1), 114 (C-5 and C-3), 67 (CH<sub>2</sub>-OAr), 52 (CH<sub>3</sub>O), 32-23 (7 CH<sub>2</sub>), 14 (CH<sub>3</sub>)


Mass spectrum:

Positive ESI  $m/z = 327.2$  [M + Na]<sup>+</sup>

30 High res. Calc. for  $\text{C}_{19}\text{H}_{28}\text{O}_3\text{Na}$ : 327.193914, Found : 327.192630.

Infrared ( $\text{cm}^{-1}$ ): 3000-2962-2925-2855-1720-1607-1511-1435-1279-1254-846

#### 4-(undec-4Z-enyloxy)benzoic acid (**34**)



5 550  $\mu$ L of 1 N sodium hydroxide solution (550  $\mu$ mol) are added portionwise to 109 mg of **33** (358  $\mu$ mol) in methanol (4 mL). The solution is refluxed for 20 hours. After evaporating off the solvent, the reaction medium is acidified with 0.5 N HCl and extracted with DCM. 102 mg of a white solid are obtained, i.e. a yield of 98%.

10  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):

12.00-11.00 (bd, 1 H,  $\text{CO}_2\text{H}$ ), 8.07 (d, 2 H,  $\text{ArH-2}$  and  $\text{ArH-6}$ ,  $J_{2,3} \approx J_{6,5}$  8.5 Hz), 6.94 (d, 2 H,  $\text{ArH-3}$  and  $\text{ArH-5}$ ), 5.42 (m, 2 H,  $\text{CH=CH}$ ), 4.03 (t, 2 H,  $\text{CH}_2\text{-OAr}$ ,  $J$  6.3 Hz), 2.26-2.03 (2 m, 4 H,  $\text{CH}_2\text{-CH=CH-CH}_2$ ), 1.88 (tt, 2 H,  $\text{ArO-CH}_2\text{-CH}_2\text{-CH}_2$ ,  $J$  6.8 Hz), 1.40-1.10 (m, 8 H, 4  $\text{CH}_2$ ), 0.89 (t, 3 H,  $\text{CH}_3$ ,  $J$  6.6 Hz)

15  $^{13}\text{C}$  NMR (62.5 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm):


172 ( $\text{C=O}$ ), 164 ( $\text{C-4}$ ), 132 ( $\text{C-2}$  and  $\text{C-6}$ ), 131-128 ( $\text{CH=CH}$ ), 121 ( $\text{C-1}$ ), 114 ( $\text{C-3}$  and  $\text{C-5}$ ), 67 ( $\text{CH}_2\text{-OAr}$ ), 32-22 (7  $\text{CH}_2$ ), 14 ( $\text{CH}_3$ )

Mass spectrum:

20 Negative ESI  $m/z$  = 289.2  $[\text{M-H}]^-$

High res. Calc. for  $\text{C}_{18}\text{H}_{25}\text{O}_3$ : 289.180370, Found : 289.178710

#### 4-(undec-4Z-enyloxy)benzoyl chloride (**35**)



25

1 mL of oxalyl chloride (11.5 mmol) and two drops of anhydrous DMF are added under argon to 101 mg of toluene-dried acid **34** (348  $\mu$ mol) dissolved in 18 mL of anhydrous DCM. The medium is stirred at room temperature for two hours and then concentrated to give 107 mg of a yellow oil, i.e. a yield of 99%.

30



<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ (ppm):

7.96 (d, 2 H, ArH-2 and ArH-6,  $J_{2,3} \approx J_{6,5}$  8.7 Hz), 6.99 (d, 2 H, ArH-3 and ArH-5), 5.43 (m, 2 H, CH=CH), 4.10 (t, 2 H, CH<sub>2</sub>-OAr,  $J$  6.3 Hz), 2.27-2.03 (2 m, 4 H, CH<sub>2</sub>-CH=CH-CH<sub>2</sub>), 1.91 (tt, 2 H, ArO-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>,  $J$  6.7 Hz), 1.35-1.12 (m, 8 H, 4 CH<sub>2</sub>), 0.89 (t, 3 H, CH<sub>3</sub>,  $J$  6.6 Hz)